










Circular dichroism in higher-order harmonic generation: Heralding topological phases and transitions in Chern insulators

Alexis Chacón ^{1,2,*}, Dasol Kim ², Wei Zhu,¹ Shane P. Kelly ^{1,3}, Alexandre Dauphin ⁴, Emilio Pisanty ⁴,
Andrew S. Maxwell,^{4,5} Antonio Picón ^{4,6}, Marcelo F. Ciappina ^{4,7,8,9}, Dong Eon Kim,² Christopher Ticknor,¹
Avadh Saxena ¹ and Maciej Lewenstein ^{4,10}

¹Center for Nonlinear Studies and Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

²Department of Physics and Center for Attosecond Science and Technology, POSTECH, 7 Pohang 37673, South Korea
and Max Planck POSTECH/KOREA Research Initiative, Pohang, 37673, South Korea

³Physics and Astronomy Department, University of California Riverside, Riverside, California 92521, USA

⁴ICFO – Institut de Ciències Fòniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels (Barcelona), Spain

⁵Department of Physics & Astronomy, University College London, Gower Street, London WC1E 6BT, United Kingdom

⁶Departamento de Química, Universidad Autónoma de Madrid, 28049 Madrid, Spain

⁷Institute of Physics of the ASCR, ELI-Beamlines project, Na Slovance 2, 182 21 Prague, Czech Republic

⁸Physics Program, Guangdong Technion – Israel Institute of Technology, Shantou, Guangdong 515063, China

⁹Technion – Israel Institute of Technology, Haifa 32000, Israel

¹⁰ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain



(Received 15 July 2020; accepted 2 September 2020; published 21 October 2020)

Topological materials are of interest to both fundamental science and advanced technologies because topological states are robust with respect to perturbations and dissipation. Experimental detection of topological invariants is thus in great demand but remains extremely challenging. Ultrafast laser-matter interactions, and in particular high-harmonic generation, meanwhile, were proposed several years ago as tools to explore the structural and dynamical properties of various matter targets. Here we show that the high-harmonic emission signal produced by a circularly polarized laser contains signatures of topological phases and transitions in the paradigmatic Haldane model. In addition to clear shifts of the overall emissivity and harmonic cutoff, the high-harmonic emission shows a unique circular dichroism, which exhibits clear changes in behavior at the topological phase boundary. Our findings pave the way to understand fundamental questions about the ultrafast electron-hole pair dynamics in topological materials via nonlinear high-harmonic generation spectroscopy.

DOI: [10.1103/PhysRevB.102.134115](https://doi.org/10.1103/PhysRevB.102.134115)

I. INTRODUCTION

The study of topological order was initiated by the discoveries of the Berezinskii-Kosterlitz-Thouless transition in 2D as well as the integer and fractional quantum Hall effects [1–3]. These materials constitute a new paradigm, since they are characterized by a global order parameter: this goes beyond the standard Landau theory of phase transitions, which uses *local* order parameters to describe materials. Topological order, due to its robustness and resistance to perturbations, has already found applications in standards and metrology (most notably, via the integer quantum Hall effect, a key ingredient of the recent revision to the SI system of units [4]), and promises numerous applications from quantum spintronics and valleytronics to quantum computing.

One question of particular interest is the possible applications of topological insulators and superconductors [5]. These systems are insulating in their bulk, but have conducting surface states protected by the topological invariant of the bulk. A considerable number of solid-state systems with these properties have been proposed in recent years, both in the context of

real topological materials [6] and in synthetic ones, employing ultracold atoms [7], photonic systems [8], and mechanical systems [9], among others. Nevertheless, new methods for the creation and physical characterization of topological phases are still being sought for real materials. Here we focus on the detection of topological phase transitions and different topological phases by the highly nonlinear optical responses of the medium—specifically, high-harmonic generation (HHG)—as a counterpart to recent proposals based on linear-optical properties [10,11].

HHG is a highly nonperturbative process in which a material medium (which may be an atom, molecule, or solid) is driven by an intense laser field, producing extreme ultraviolet (XUV) photons as harmonics of the driving frequency. The XUV emission can carry information about the structure and dynamics of the medium, as first proposed for molecules by Itatani *et al.* [12] (for a recent review of various processes in strong laser fields, see Ref. [13]). In HHG, an ultrashort (5–50 fs) intense midinfrared (MIR) laser pulse causes partial ionization of an electron in an atom or molecule; the resulting electronic wave packet is accelerated in the laser field, returns to and recombines with the parent ion, producing high-order harmonics [14,15]. The efficiency of this process depends directly on the atomic or molecular orbital

*achacon@postech.ac.kr

that the electron leaves and recombines with. The electronic and molecular structure of the target can similarly be probed by the rescattered electron (without recombination) via laser-induced electron diffraction [16,17].

In the last few years, the subject of HHG from solid-state targets has attracted considerable attention [18–20]. In particular, Vampa *et al.* investigated the role of intraband and interband currents in ZnO experimentally, employing HHG to characterize structural information such as the energy dispersions [21–23] from their nontrivial coupling [24], where interband dynamics governed the emission. However, depending on the material structure, Golde *et al.* [24] also showed that the intraband mechanism might govern over the interband. Additionally, real-space approaches that, in some regimes, discard the band-structure picture altogether have recently been developed [25]. The latter show that inter- and intraband mechanisms cannot trivially be disentangled, and their natures and relationship remain a matter of intense discussion in ultrafast science.

Generally, existing studies have dealt with standard materials, where the topology does not play a role. Nevertheless, Berry-phase effects have been explored in some topologically trivial materials: This includes experimental studies of HHG in atomically thin semiconductors [26] and in quasi-2D models [27], which show the sensitivity of harmonic emission to symmetry breaking (specifically, the breaking of inversion symmetry (IS) in monolayer MoS₂ and α quartz, via the appearance of even harmonics), as well as the effects of Berry curvature on the HHG spectrum of a time-reversal-invariant material [28]. On a broader outlook, recent works have focused on HHG in graphene [29–32], monolayer transition-metal dichalcogenides [33,34], and strained MoS₂ [35,36], as well as bilayer graphene [37]. HHG from strongly correlated materials has also been investigated recently, with work probing dynamical Mott-insulator transitions [38] as well as strongly correlated spin systems [39].

Very recent work, however, has started to consider more directly the question of whether HHG is sensitive to topological order, starting from the contributions of edge states in a 1D-chain model of the Su-Schrieffer-Heeger (SSH) system [40], which can significantly influence the harmonic emission [41–43]. In particular, recent work has also shown that the topological transitions in the Haldane model (HM)—a paradigmatic Chern insulator—leave explicit traces in the helicity of the emitted harmonics [44]. Further work has also probed HHG in carbon nanotubes [45], few-layer [46], and 3D [47] topological insulators, and in materials with nodal topology [48].

Interestingly, Tran *et al.* [11] considered Chern insulators in an ultracold atom system in circularly shaken lattices, corresponding to impinging a circularly polarized driver. They demonstrated that the depletion rates corresponding to left- and right-polarized drivers show dependence on the orientation of the circular shaking. This *circular dichroism* (CD) was then directly related to the topological invariant, the so-called Chern number. The latter was recently measured in a cold atom experiment [49]. In this paper, we show that the CD produced with HHG can be used to herald the topological phase transition in a 2D topological Chern insulator material, analogous to its role in magnetic materials [50]. We

demonstrate this in detail for the paradigmatic HM, illustrated in Fig. 1 [51]. We derive, verify, and apply the theory of HHG driven by both linearly and circularly polarized light in a two-band model, fully including the effects of the Berry curvature and its associated connection. We characterize and analyze the HHG spectrum, via both the intra- and interband contributions, and using both numerical and analytical approaches.

Moreover, we develop a theoretical formalism which is gauge invariant with respect to both the electromagnetic and the Bloch wave-function gauges [52]. In topological phases, no continuous single-gauge charts exist [53] which would allow a correct and regular description of dipole transition matrix elements over the entire Brillouin zone (BZ): for every gauge, there is at least one point in k space where the dipole moment (and with it the Berry connection) has a discontinuity. To deal with this inevitable discontinuity, we develop a multigauge approach using separate Bloch-gauge charts for different regions of the BZ. We also perform a crystal-momentum saddle-point analysis of the obtained expressions for the interband and intraband currents. This allows us to derive expressions for the semiclassical electron-hole trajectories—including the effect of the band topology and Berry curvature and connection—and also to preserve gauge invariance (in contrast to, e.g., Ref. [54]) accounting explicitly for the phase of the transition dipole moments.

Our theoretical approach provides a complete model which predicts the following: (i) reported features of experiments for both linearly and circularly polarized driving lasers [26,27,55] and novel behaviors for topologically non trivial systems; (ii) that HHG is extremely sensitive to IS and, in addition, to breaking time-reversal symmetry (TRS); and (iii) the HHG spectrum intensity depends on the topological phases, and the CD of high harmonic orders (HOs) is sensitive to the crossing of a topological phase transition boundary. Our work is mainly focused on the HM but we expect that the broad features are generally applicable to *Chern insulators*.

After this Introduction, we review the HM in Sec. II. Next, in Sec. III, we describe in detail how to handle the singularities of the dipole transition moment which are required to appear in topologically nontrivial phases. In Sec. IV, we discuss the semiconductor Bloch equations (SBEs) and methods of their treatment, as well as our observables, the inter- and intraband charge currents. Section V is devoted to the discussion of the Keldysh approximation, as well as a quasiclassical analysis based on the saddle point approximation applied to the gauge-invariant action, from which we can analyze the effect of the topology on the electron-hole trajectories. In Sec. VI, we begin to examine applications of our theory and consider the first HHG spectra in practice as a benchmark for our theory against known results for α -quartz. We go on, in Sec. VII, to discuss the question whether HHG can provide signatures of topological phases in any sense. We argue that the answer to this question is positive, and this is particularly spectacular when we consider CD, which we discuss in detail in Sec. VIII. Additional support for our results can be gathered by analyzing the effects of dephasing on HHG spectra: The harmonics that exhibit strong dichroism are particularly robust with respect to dephasing, which we explore in Sec. IX. Finally, we conclude and present an outlook in Sec. X.




FIG. 1. Phase diagram and band structures of the Haldane model. (a) Phase diagram in the plane $(\phi_0, M_0/t_2)$, with the different phases labeled by their Chern number C ; the green line shows the boundary of the topological phase transition. The band-structure diagrams correspond to the points X, Y, Z, and Q as marked in (a), with the point Q at the phase transition showing the gapless Dirac cone at the K point. (Y) and (Z) depict the band structure when the conduction-band topological invariants are $C = \pm 1$ at the phase points $\phi_0 = \pm\pi/2$ and $M_0 = 2.54t_2$. (Y) shows how the midinfrared laser-source oscillations (red-solid line) drive the topological material; this can be with both linear and circular polarizations. We also depict a physical cartoon of the electron-hole pair dynamics driven by a linearly polarized laser, i.e., creation, propagation, and annihilation or recombination by the black-dashed lines with arrows, and finally the subsequent harmonic emission (violet oscillations). The dashed lines in (a) indicate the cuts used for the parameter scans below. Panels (b) and (c) show the Brillouin zone and the real-space lattice of the Haldane model with the couplings in use.

II. THE HALDANE MODEL

The HM [51], originally introduced as a toy model, represents the first example of an anomalous quantum Hall effect [56], and it captures the essential features of a number of materials [57,58]—in particular, the quantized transverse (spin) conductivities of the quantum spin Hall effect [59], where the transport takes place along protected edge states. Moreover, this model remains solvable and implementable in quantum simulators [60], which makes it a flexible tool for understanding a wide range of phenomena. Specifically, the HM describes a tight-binding Hamiltonian of spinless fermions on a 2D hexagonal lattice with a real nearest-neighbor (NN) hopping t_1 , an on-site staggering potential M_0 , and a complex next-to-nearest-neighbor (NNN) hopping $t_2 e^{i\phi_0}$. The HM belongs to the class of Chern insulators [61] and is characterized by a topological invariant, i.e., the Chern number [see Eq. (11)]. In this section, we provide the details of the Hamiltonian, its eigenvalues and eigenvectors, its Berry connection, the Berry curvature, and Chern number that derive from it, and the transition dipole moments, as well as the relationships between these quantities.

A. Hamiltonian for the Haldane model

The HM is described by a tight-binding Hamiltonian of spinless fermions with a real NN hopping t_1 , a complex NNN hopping $t_2 e^{i\phi_0}$, and an on-site staggering potential M_0 . The

Hamiltonian in momentum space reads

$$H_0(\mathbf{k}) = B_{0,\mathbf{k}} \sigma_0 + \mathbf{B}_{\mathbf{k}} \cdot \boldsymbol{\sigma}, \quad (1)$$

where the set of $B_{0,\mathbf{k}}$ and $\mathbf{B}_{\mathbf{k}} = \{B_{1,\mathbf{k}}, B_{2,\mathbf{k}}, B_{3,\mathbf{k}}\}$ is known as a pseudomagnetic field describing the tight-binding components of the Hamiltonian in momentum space, which read

$$B_{0,\mathbf{k}} = 2t_2 \cos \phi_0 \sum_{i=1}^3 \cos(\mathbf{k} \cdot \mathbf{b}_i), \quad (2a)$$

$$B_{1,\mathbf{k}} = t_1 \sum_{i=1}^3 \cos(\mathbf{k} \cdot \mathbf{a}_i), \quad (2b)$$

$$B_{2,\mathbf{k}} = t_1 \sum_{i=1}^3 \sin(\mathbf{k} \cdot \mathbf{a}_i), \quad (2c)$$

$$B_{3,\mathbf{k}} = M_0 - 2t_2 \sin \phi_0 \sum_{i=1}^3 \sin(\mathbf{k} \cdot \mathbf{b}_i). \quad (2d)$$

Here the \mathbf{a}_i are the NN vectors of the lattice [from atom A to B, see Fig. 1(c) for more details], $\mathbf{a}_1 = (0, a_0)$, $\mathbf{a}_2 = \frac{1}{2}(-\sqrt{3}, -1)a_0$ and $\mathbf{a}_3 = \frac{1}{2}(\sqrt{3}, -1)a_0$, and the \mathbf{b}_i are the NNN vectors (from atom A to A, or B to B) given by $\mathbf{b}_1 = (\sqrt{3}, 0)a_0$, $\mathbf{b}_2 = \frac{1}{2}(-\sqrt{3}, +3)a_0$ and $\mathbf{b}_3 = \frac{1}{2}(-\sqrt{3}, -3)a_0$, with a_0 denoting the distance between the atoms A and B. In Eq. (1), moreover, σ_0 is the identity matrix and $\boldsymbol{\sigma} = \{\sigma_1, \sigma_2, \sigma_3\}$ is the vector of Pauli matrices, which is taken on a reference frame such that the eigenstates of σ_3 are Bloch

waves that are localized to sites A and B, respectively, within each unit cell. The symmetries of the model are obtained as follows: (i) a nonzero magnetic flux ϕ_0 breaks the TRS and (ii) a nonzero staggering potential M_0 breaks the IS.

B. Eigenvalues and eigenvectors

As the Hamiltonian is described by a 2×2 matrix in the Bloch basis, the eigenvalues and the eigenvectors of the Hamiltonian can be found analytically, using standard tools: the Hamiltonian is essentially a Pauli matrix along the $\hat{\mathbf{B}}(\mathbf{k}) := \mathbf{B}_k/|\mathbf{B}_k|$ direction in the same space as the σ Pauli vector and its eigenvectors are the corresponding spin states as usual. The energy spectrum thus reads

$$\varepsilon_{c/v}(\mathbf{k}) = B_{0,\mathbf{k}} \pm |\mathbf{B}_k|, \quad (3)$$

with the upper (lower) sign corresponding to the conduction (valence) band. We also define, for future use, the polar coordinates ϕ_k and θ_k of $\hat{\mathbf{B}}(\mathbf{k})$, as $\tan \phi_k := \frac{\hat{B}_{2,\mathbf{k}}}{\hat{B}_{1,\mathbf{k}}}$ and $\cos \theta_k := \hat{B}_{3,\mathbf{k}}$.

The eigenvectors of H_0 at crystal momentum \mathbf{k} are the Bloch waves

$$|\Phi_{m,\mathbf{k}}\rangle = e^{i\mathbf{k}\cdot\mathbf{r}}|u_{m,\mathbf{k}}\rangle, \quad (4)$$

where $|u_{m,\mathbf{k}}\rangle$ are the lattice-periodic functions which are given, in the basis of A- and B-Bloch waves that the Hamiltonian Eq. (1) acts in, by

$$|u_{+,\mathbf{k}}\rangle = \frac{1}{N_{+,\mathbf{k}}} \begin{pmatrix} B_{3,\mathbf{k}} + |\mathbf{B}_k| \\ B_{1,\mathbf{k}} + iB_{2,\mathbf{k}} \end{pmatrix} = \begin{pmatrix} e^{-i\phi_k/2} \cos \frac{\theta_k}{2} \\ e^{+i\phi_k/2} \sin \frac{\theta_k}{2} \end{pmatrix}, \quad (5)$$

$$|u_{-,\mathbf{k}}\rangle = \frac{1}{N_{-,\mathbf{k}}} \begin{pmatrix} iB_{2,\mathbf{k}} - B_{1,\mathbf{k}} \\ B_{3,\mathbf{k}} + |\mathbf{B}_k| \end{pmatrix} = \begin{pmatrix} e^{-i\phi_k/2} \sin \frac{\theta_k}{2} \\ -e^{+i\phi_k/2} \cos \frac{\theta_k}{2} \end{pmatrix}, \quad (6)$$

with $N_{\pm,\mathbf{k}}$ a suitable normalization constant. As usual, these eigenstates are only defined up to a (possibly \mathbf{k} -dependent) phase. Changes to this phase are the Bloch gauge transformations that we will examine in depth in Sec. III. A gauge transformation of the wave function Eqs. (5) and (6) can be performed by a unitary transformation and would lead to a new vector $\mathbf{B}'_k = \{B'_{1,\mathbf{k}}, B'_{2,\mathbf{k}}, B'_{3,\mathbf{k}}\}$ (for example, $\mathbf{B}' = \{B_2, B_3, B_1\}$). Such a transformation keeps invariant the energy dispersions and Berry curvatures but modifies the Berry connection and the transition dipole moments. This gauge control will be implemented numerically to manipulate the discontinuities and singularities of the dipole matrix element and Berry connections in the BZ of topological phases.

C. Berry connection, Berry curvature, and Chern number

To calculate the radiation-interaction properties of these eigenstates, we require the dipole transition matrix elements between them. These can be singular functions of \mathbf{k} [62,63] and read

$$\langle \Phi_{m',\mathbf{k}'} | \mathbf{x} | \Phi_{m,\mathbf{k}} \rangle = -i \nabla_{\mathbf{k}} (\delta_{m'm} \delta(\mathbf{k} - \mathbf{k}')) + \delta(\mathbf{k} - \mathbf{k}') \mathbf{d}_{m'm}(\mathbf{k}), \quad (7)$$

where

$$\mathbf{d}_{m'm}(\mathbf{k}) = i \langle u_{m',\mathbf{k}} | \nabla_{\mathbf{k}} | u_{m,\mathbf{k}} \rangle \quad (8)$$

is a regular function which encodes the momentum gradient of the periodic part of the Bloch functions. In Sec. III below, we will examine in detail the off-diagonal matrix elements $\mathbf{d}_{cv}(\mathbf{k})$, which must always exhibit singularities when the material is in the topological phase, but for now we focus on the diagonal elements. In 1D systems with trivial topology, the diagonal elements of the transition dipoles can always be made to vanish but in higher dimensionality this is impossible: these diagonal elements give the Berry connection of the band,

$$\begin{aligned} \xi_m(\mathbf{k}) &= \mathbf{d}_{mm}(\mathbf{k}) \\ &= i \langle u_{m,\mathbf{k}} | \nabla_{\mathbf{k}} | u_{m,\mathbf{k}} \rangle, \end{aligned} \quad (9)$$

which is responsible for the parallel transport of wave-function phase around the band. This parallel transport is measured by the Berry curvature, given by the gauge-invariant curl

$$\Omega_m(\mathbf{k}) = \nabla_{\mathbf{k}} \times \xi_m(\mathbf{k}) \quad (10)$$

of the connection, whose integral over the entire band,

$$C_m := \frac{1}{2\pi} \int_{\text{BZ}} \Omega_m(\mathbf{k}) \cdot d^2\mathbf{k}, \quad (11)$$

is the topological invariant of the system, known as the Chern number.

For the specific case of the HM eigenstates with the gauge fixed by Eqs. (5) and (6), the Berry connection is given explicitly by

$$\xi_{v/c}(\mathbf{k}) = \mp \frac{1}{2} (\cos \theta_k) (\nabla_{\mathbf{k}} \phi_k), \quad (12)$$

so the Berry curvature reads

$$\Omega_{v/c}(\mathbf{k}) = \nabla_{\mathbf{k}} \times \xi_{v/c}(\mathbf{k}) = \mp \frac{1}{2} (\nabla_{\mathbf{k}} \cos \theta_k) \times (\nabla_{\mathbf{k}} \phi_k), \quad (13)$$

and the Chern number results are

$$C_{c/v} = \mp \frac{1}{4\pi} \int_{\text{BZ}} d^2\mathbf{k} \cdot [\sin \theta_k (\nabla_{\mathbf{k}} \theta_k) \times (\nabla_{\mathbf{k}} \phi_k)]. \quad (14)$$

In the HM, the Chern number of the conduction band $C_c = C$ can take the values of -1 , 0 , and $+1$ depending on M_0 and ϕ_0 (see Fig. 1). The trivial phase has a zero Chern number and the two topological phases are characterized by Chern numbers $C = \pm 1$. Figure 1(a) depicts the phase diagram in terms of ϕ_0 and M_0 , showing three distinct phases: a trivial phase with Chern number $C = 0$ and two different topological phases, with Chern numbers $C = \pm 1$. At the boundary between the phases, the band gap closes, as shown for the point (Q).

D. Dipole moment and the Berry curvature in the Haldane model

We now turn to the dipole matrix element, as initially defined in Eq. (8), and which is here given by

$$\mathbf{d}_{cv}(\mathbf{k}) = \frac{1}{2} [(\sin \theta_k) (\nabla_{\mathbf{k}} \phi_k) + i \nabla_{\mathbf{k}} \theta_k]. \quad (15)$$

On the one hand, we find that the cross product of the HM dipole Eq. (15) yields

$$\text{Im} (d_{cv}^x d_{cv}^{y*}) = \frac{1}{4} \sin \theta_k [\partial_{k_y} \theta_k \partial_{k_x} \phi_k - \partial_{k_x} \theta_k \partial_{k_y} \phi_k]. \quad (16)$$

Expanding the Berry curvature in the HM, given by Eq. (13), one obtains

$$\begin{aligned} \Omega_{v/c}(\mathbf{k}) &= \mp \frac{1}{2} \sin \theta_k (\nabla_{\mathbf{k}} \theta_k \times \nabla_{\mathbf{k}} \phi_k) \\ &= \mp \hat{\mathbf{z}} \frac{1}{2} \sin \theta_k [\partial_{k_y} \theta_k \partial_{k_x} \phi_k - \partial_{k_x} \theta_k \partial_{k_y} \phi_k]. \end{aligned} \quad (17)$$




FIG. 2. (a)–(d) Dipole transition matrix elements over the Brillouin zone for trivial and topological phases for different gauges. We show the x component in a color map ranging from dark blue to red, with white at zero. The hexagon shows the first BZ, with the K (K') points denoted by the white (yellow) circles. The insets in (c), (d) show cuts through the singularities: (c) along k_y , at fixed $k_x = -1.85$ (black), -1.80 (green), -1.75 (blue), and -1.70 a.u. (red), and (d) in vertical cuts at fixed $k_y = -0.55$ (black), -0.50 (green), -0.45 (blue), and -0.40 a.u. (red). The cyan circles highlight the discontinuities in the dipoles. (e)–(h) The resulting HHG spectra under an LCP laser driver, for gauges A and B (e), (g) and our variable-gauge method (f), (h), for both phases. We set $t_1 = 0.075$ a.u., $t_2 = t_1/3$, $M_0 = 0.0635$ a.u. = $2.54t_2$ and $a_0 = 1$ Å, giving a band gap of $E_g = 3.0$ eV, and we use a magnetic flux of (a), (b) $\phi_0 = 0.06$ rad, giving $C = 0$, and (c), (d) $\phi_0 = 1.16$ rad, giving $C = +1$. For the HHG spectra, we use the pulse Eq. (25) with $E_0 = 0.0045$ a.u., $\omega_0 = 0.014$ a.u. = 0.38 eV (corresponding to $\lambda = 3.2$ μm), and with a Gaussian envelope lasting $N_c = 14$ cycles at FWHM; we use a dephasing time of $T_2 = 220$ a.u. We integrate over a momentum grid of $N_x = 873$ by $N_y = 511$ points using a two-dimensional Simpson rule, with boundaries at $k_{x,\text{max}} = \frac{2\pi}{\sqrt{3}a_0}$ and $k_{y,\text{max}} = \frac{2\pi}{3a_0}$; for the SBE time integration our time step is $dt = 0.1$ a.u.

We conclude, thus, that

$$\Omega_{v/c} = \mp 2 \text{Im} [d_{cv}^{(x)} d_{cv}^{*(y)}], \quad (18)$$

which demonstrates the close relationship between dipole matrix elements and the Berry curvature in the HM. This confirmation of the relation of the dipole product with the Berry curvature is extremely important, since this leads to a direct connection of the interband transition current of Eq. (36) and the topological invariant for this model.

III. GAUGE-INDUCED DISCONTINUITIES IN THE TRANSITION DIPOLES

In general, it is desirable for the phase in front of the eigenstates in Eqs. (5) and (6), together with the Berry connections and the dipole matrix elements that derive from it, to be continuous over the entire Brillouin zone (BZ). However, this is not always possible: Indeed, this is the *core distinction* between the topologically trivial and nontrivial phases. In the nontrivial phase, it is provably impossible to find a Bloch wave function gauge that will work smoothly for all momenta in the BZ [53].

In the trivial phase, on the other hand, globally smooth gauges are viable [52]—but they are not guaranteed [63], either, so one must always be prepared to handle discontinuities and singularities in these quantities. Moreover, as the SBE directly incorporate these two quantities [see below Eqs. (23) and (24)], the numerical calculations of the currents can inherit those artificial singularities, leading to numerical instabilities and noise in the harmonic plateau

or cutoff. Here we show an example of how this numerical instability arises and the method we use to solve it. This method relies on the control of the localization of the dipole discontinuity in the BZ. Such control is realized by choosing different gauges for different regions (in the language of differential geometry, for different charts [64]) of the BZ.

Figures 2(a) and 2(b) show the real part of the x component of the dipole matrix element, $d_{cv}^{(x)}(\mathbf{k})$, of a trivial topological phase, taken in two different gauges: A, where $\mathbf{B}'_{\mathbf{k},A} = (B_{2,\mathbf{k}}, B_{3,\mathbf{k}}, B_{1,\mathbf{k}})$, and B, where $\mathbf{B}'_{\mathbf{k},B} = (B_{3,\mathbf{k}}, B_{1,\mathbf{k}}, B_{2,\mathbf{k}})$. As can be seen in the figure, both gauges exhibit substantially different dipoles, but their corresponding high-harmonic emissions—shown in Fig. 2(e)—are absolutely the same, i.e., *the harmonic emission is gauge invariant*, as expected. This is a simple and obvious test, but it demonstrates a powerful mathematical property that will help us to circumvent the numerical noise, which can compromise our (or any) interpretations of the harmonic emission from topological phases, if the singularities are not well handled or numerically integrated. Here, we find a simple method to solve this issue.

Figures 2(c), 2(d), and 2(g) show similar results for a topologically nontrivial phase. Most prominently, in the dipole maps of Figs. 2(c) and 2(d), we notice pronounced discontinuities in the dipole for both gauges, which are highlighted by the cyan circles (see inset plots). Second, the HHG calculations for gauges A and B, shown in Fig. 2(f), show numerical noise around the plateau at HOs $> 5^{\text{th}}$ and, more troublingly, a lack of a well-defined cutoff. These features signal a numerical instability of the calculation, which is independent

of the numerical convergence of our fifth-order Runge–Kutta method.

To solve this problem, we use the fact that the k -space discontinuities in the dipole can be shifted within the BZ [as shown in Figs. 2(c) and 2(d)] by choosing different gauges: the existence of singularities is inevitable [53] but their location in \mathbf{k} -space is not fixed and it can be moved by switching phase conventions. Therefore, we use different gauges to calculate the SBE [see Eqs. (23) and (24)] evolution in different patches of the BZ, and then coherently combine the currents at the end of the calculation. Thus, in our variable-gauge technique, we apply the two gauges described above, A and B, for the different BZ regions where each gauge is regular, using the following three criteria:

(i) Gauge A for the BZ region which obeys the constraints $-k_{y,\max} \leq k_y \leq \epsilon_{y,1}$, where $\epsilon_{y,1} = -0.35$ a.u. is marked with the horizontal green dashed line in Fig. 2(d).

(ii) Since all physical quantities are periodic within the whole BZ, the regions Q1, Q2, and Q3 [marked by green boxes in Fig. 2(c)] are physically equivalent, we use gauge A shifted to the Q3 region when k_y satisfies $\epsilon_{y,1} < k_y \leq \epsilon_{y,2}$, where $\epsilon_{y,2} = 0.02$ a.u.

(iii) We use gauge B for momenta which satisfy $\epsilon_{y,2} < k_y \leq k_{y,\max}$.

We compute HHG with this variable-gauge method for both the trivial and topological phases. For the trivial phase we show the resulting harmonic spectrum in Fig. 2(h), where the red dashed line shows the variable-gauge results; we find very good agreement between this approach and the fixed-gauge reference (using gauge B), which works well for the trivial phase, as discussed above, and thus serves as a benchmark for the variable-gauge approach. Silva *et al.* have developed an interesting alternative to circumvent the dipole matrix elements singularities in trivial and topological materials by integrating numerically the SBEs or the reduced density matrix, namely, the maximally localized Wannier-approach (MLW) framework [65]. Our *variable gauge approach* was compared to the MLW. This variable-gauge Bloch wave function method perfectly agrees with MLW too.

With this in place, we can confidently apply the same technique to the calculation of HHG for the topological phase, the results of which are shown in the red line of Fig. 2(h). The HHG spectrum for this topological phase shows a clean plateau with well-defined harmonics and a clear cutoff region. Moreover, there is a clear numerical threshold, in comparison to the fixed-gauge method using gauge B. Our numerical threshold is fixed at $I_{\text{HHG}} \approx 10^{-16}$ arb. units, but we have found that this threshold can be reduced up to 10^{-25} arb. units for the HHG yield, depending on the parameters used for the BZ grid and the time integration.

That said, it is important to note that, while our variable-gauge method can map the plateau and cutoff very well for topological phases, our method finds an undefined harmonic cutoff, as shown in Fig. 6 below. This occurs at the critical point of the HM, where the band gap closes, at the topological phase transition. This behavior is expected, since the closing of the band gap creates an undetermined dipole matrix element at the Dirac cone, where the population is most easily transferred to the conduction band.

IV. SEMICONDUCTOR BLOCH EQUATIONS

We introduce and examine the equations governing the microscopic time-dependent response of the solid due to the laser-matter interaction, the SBEs [66]. We follow the approach of Refs. [18,19], generalizing it to include the effects of possible nontrivial topology.

A. Semiconductor Bloch equations

If we write down the electronic-crystalline wave function in the form

$$|\Psi(t)\rangle = \sum_m \int_{\text{BZ}} d\mathbf{k} a_m(\mathbf{k}, t) |\Phi_{m,\mathbf{k}}\rangle, \quad (19)$$

the time evolution of the Bloch-space probability amplitudes $a_m(\mathbf{k}, t)$, as determined by the Schrödinger equation, can be transformed into the form [62]

$$i\dot{a}_m(\mathbf{k}', t) = \varepsilon_m(\mathbf{k}') a_m(\mathbf{k}', t) + \mathbf{E}(t) \cdot \sum_{m'} \int_{\text{BZ}} \langle \Phi_{m,\mathbf{k}'} | \mathbf{x} | \Phi_{m',\mathbf{k}} \rangle a_{m'}(\mathbf{k}', t). \quad (20)$$

Here $\varepsilon_m(\mathbf{k})$ denotes the energy dispersion for the valence/conduction band $m = v, c$, and m' also ranges over both bands. We then compute the second and third terms on the right-hand side of the above equation by using Eq. (7), and we find

$$\begin{aligned} \dot{a}_m(\mathbf{k}, t) = & -i[\varepsilon_m(\mathbf{k}) + \mathbf{E}(t) \cdot \boldsymbol{\xi}_m(\mathbf{k}) - i\mathbf{E}(t) \cdot \nabla_{\mathbf{k}}] a_m(\mathbf{k}, t) \\ & - i\mathbf{E}(t) \cdot \sum_{m' \neq m} \mathbf{d}_{mm'}(\mathbf{k}) a_{m'}(\mathbf{k}, t). \end{aligned} \quad (21)$$

This differential equation mixes time derivatives with the momentum gradient $\nabla_{\mathbf{k}}$, which also occurs for strong field approximation (SFA) treatment of atoms [13]. Thus, we apply the same transformation, shifting the momentum over time to $\mathbf{K} = \mathbf{k} - \mathbf{A}(t)$ —the canonical crystal quasi-momentum, which is a constant of motion—giving rise to a shifted Brillouin zone, denoted BZ . More formally, by applying the substitution $a_m(\mathbf{k}, t) = e^{i\mathbf{A}(t) \cdot \nabla_{\mathbf{k}}} b_m(\mathbf{K}, t)$, one finds that

$$\begin{aligned} \dot{b}_m(\mathbf{K}, t) = & -i[\varepsilon_m(\mathbf{K} + \mathbf{A}(t)) + \mathbf{E}(t) \cdot \boldsymbol{\xi}_m(\mathbf{K} + \mathbf{A}(t))] b_m(\mathbf{K}, t) \\ & - i\mathbf{E}(t) \cdot \sum_{m' \neq m} \mathbf{d}_{mm'}(\mathbf{K} + \mathbf{A}(t)) b_{m'}(\mathbf{K}, t). \end{aligned} \quad (22)$$

We then transform the transition amplitude b_m to the density matrix operator $\hat{\rho}$, i.e., the population $n_m = \rho_{mm}$ and coherence $\pi = \rho_{cv}$, given explicitly by $n_m(\mathbf{K}, t) = b_m^*(\mathbf{K}, t) b_m(\mathbf{K}, t)$, and $\pi(\mathbf{K}, t) = b_c(\mathbf{K}, t) b_v^*(\mathbf{K}, t)$. Thus, this leads to the SBEs, which describe the laser-electron interaction in the lattice [24] in terms of the band population and coherence:

$$\dot{n}_m(\mathbf{K}, t) = i(-1)^m \mathbf{E}(t) \cdot \mathbf{d}_{cv}^*(\mathbf{K} + \mathbf{A}(t)) \pi(\mathbf{K}, t) + \text{c.c.}, \quad (23)$$

$$\begin{aligned} \dot{\pi}(\mathbf{K}, t) = & -i \left[\varepsilon_g(\mathbf{K} + \mathbf{A}(t)) + \mathbf{E}(t) \cdot \boldsymbol{\xi}_g(\mathbf{K} + \mathbf{A}(t)) - i \frac{1}{T_2} \right] \pi(\mathbf{K}, t) \\ & - i\mathbf{E}(t) \cdot \mathbf{d}_{cv}(\mathbf{K} + \mathbf{A}(t)) w(\mathbf{K}, t). \end{aligned} \quad (24)$$

Here $\varepsilon_g(\mathbf{k}) = \varepsilon_c(\mathbf{k}) - \varepsilon_v(\mathbf{k})$ is the energy gap between the conduction and valence bands, as a function of the crystal momentum \mathbf{k} , $\xi_g(\mathbf{k}) = \xi_c(\mathbf{k}) - \xi_v(\mathbf{k})$ is the difference in the Berry connection between the conduction and valence bands, and the sign $(-1)^m$ takes the values $(-1)^c = 1$ and $(-1)^v = -1$. We also introduce a phenomenological dephasing term written in terms of the dephasing time T_2 [19]. Finally, $w(\mathbf{k}, t) = n_v(\mathbf{k}, t) - n_c(\mathbf{k}, t)$ denotes the momentum and time dependence of the population difference between the valence- and conduction-band or inversion population.

In our calculations, we describe the driving laser using the vector potential

$$\mathbf{A}(t) = \frac{E_0}{\omega_0} f(t) \left(\frac{1}{\sqrt{1+\epsilon^2}} \cos(\omega_0(t-t_0)) \mathbf{e}_x + \frac{\epsilon}{\sqrt{1+\epsilon^2}} \sin(\omega_0(t-t_0)) \mathbf{e}_y \right), \quad (25)$$

where E_0 is the laser field strength, ω_0 is the central photon frequency, and $f(t)$ is the laser-pulse envelope, given by $f(t) = \exp[-(t-t_0)^2/(2\sigma^2)]$, where σ is the pulse width. Here ϵ denotes the ellipticity of the laser, given by $\epsilon = -1$ ($+1$) for right-handed (left-handed) circularly polarized laser fields, abbreviated RCP (LCP), respectively, and by $\epsilon = 0$ for linearly polarized laser drivers. The electric field is given by $\mathbf{E}(t) = -\partial_t \mathbf{A}(t)$.

We numerically solve the SBEs described by Eqs. (23) and (24) with a 5th-order Runge-Kutta scheme implemented in C++, using the Message Passing Interface (MPI) for parallelization. Additionally, we validate our numerical findings in C++ by comparing our numerical outcomes with the numerical solution obtained by the built-in solvers on Wolfram Mathematica. We find very good agreement between the C++ and Mathematica results.

B. Microscopic currents

The HHG spectra are obtained by Fourier analyzing microscopic charge currents induced by the driving laser pulse. Here we derive the total microscopic current $\mathbf{J}(t) = \mathbf{J}_{\text{ra}}(t) + \mathbf{J}_{\text{er}}(t)$, by focusing on its two components: the intraband $\mathbf{J}_{\text{ra}}(t)$ and the interband $\mathbf{J}_{\text{er}}(t)$ currents [18,19].

1. Interband current

We start by writing the interband current $\mathbf{J}_{\text{er}}(t) = \frac{d}{dt} \mathbf{P}_{\text{er}}(t)$ in terms of the polarization

$$\begin{aligned} \mathbf{P}_{\text{er}}(t) &\equiv e \langle \Psi(t) | \mathbf{x} | \Psi(t) \rangle \\ &= e \int_{\text{BZ}} d\mathbf{k}' \int_{\text{BZ}} d\mathbf{k} a_c(\mathbf{k}', t) a_v^*(\mathbf{k}, t) \\ &\quad \times \langle \Phi_{v,\mathbf{k}'} | \mathbf{x} | \Phi_{c,\mathbf{k}} \rangle + \text{c.c.}, \end{aligned} \quad (26)$$

where $e = -1$ a.u. is the electron charge. Next, using the identities above, the current simplifies to

$$\mathbf{J}_{\text{er}}(t) = e \frac{d}{dt} \int_{\text{BZ}} d^2\mathbf{K} \mathbf{d}_{cv}^*(\mathbf{K} + \mathbf{A}(t)) \pi(\mathbf{K}, t) d^2\mathbf{K} + \text{c.c.}, \quad (27)$$

indexed by the constant canonical crystal quasimomentum \mathbf{K} . This result coincides with the ones found by Vampa *et al.* in Ref. [67].

2. Intraband current

We write the intraband current, defined as $\mathbf{J}_{\text{ra}}(t) = e \sum_m \langle \Psi(t) | \mathbf{v}_m | \Psi(t) \rangle$, in terms of the intraband velocity operator $\mathbf{v}_m = -i[\mathbf{x}_m, H_0] - i[\mathbf{x}_m, \mathbf{x}_m \cdot \mathbf{E}(t)]$. We use the electromagnetic length gauge with the single-active-electron Hamiltonian and the dipole approximation of the laser-crystal system, i.e., $H(t) = H_0 + \mathbf{x} \cdot \mathbf{E}(t)$. For simplicity, our derivation is focused on the conduction band $m = c$ of the intraband current, which reads

$$\begin{aligned} \mathbf{J}_{\text{ra}}(t) &= e \int d\mathbf{k}' \int d\mathbf{k} a_{c,\mathbf{k}'}^*(t) a_{c,\mathbf{k}}(t) \langle \Phi_{c,\mathbf{k}'} | \mathbf{x}_c | \Phi_{c,\mathbf{k}} \rangle \varepsilon_c(\mathbf{k}) \\ &\quad + (c \rightarrow v) \\ &= e \int d\mathbf{k}' [n_c(\mathbf{k}', t) \mathbf{v}_c(\mathbf{k}') + n_v(\mathbf{k}', t) \mathbf{v}_v(\mathbf{k}')]. \end{aligned} \quad (28)$$

Here, $\mathbf{v}_c(\mathbf{k}) = \mathbf{v}_{\text{gr},c}(\mathbf{k}) + \mathbf{v}_{a,c}(\mathbf{k})$ is written in terms of the group velocity $\mathbf{v}_{\text{gr},c}(\mathbf{k}) = \nabla_{\mathbf{k}} \varepsilon_c(\mathbf{k})$, and the anomalous velocity $\mathbf{v}_{a,c}(\mathbf{k}) = -\mathbf{E}(t) \times \boldsymbol{\Omega}_c(\mathbf{k})$, where $\boldsymbol{\Omega}_m(\mathbf{k}) = \nabla_{\mathbf{k}} \times \boldsymbol{\xi}_m(\mathbf{k})$ is the Berry curvature. The anomalous-velocity term comes from considering that the intraband component $x_m^{(j)}$ of the position operator does not commute with $x_m^{(i)}$ component, with $i, j = x, y$ for 2D [62]. This intraband current also can be expressed on the shifted Brillouin zone $\bar{\text{BZ}}$, as

$$\mathbf{J}_{\text{ra}}(t) = e \sum_m \int_{\bar{\text{BZ}}} d^2\mathbf{K} \mathbf{v}_m(\mathbf{K} + \mathbf{A}(t)) n_m(\mathbf{K}, t) d^2\mathbf{K}. \quad (29)$$

To calculate the HHG spectrum from a crystal, we have to evaluate Eqs. (27) and (29). Hence, our main task in the following will be to compute the coherence $\pi(\mathbf{k}, t)$ and the occupations $n_m(\mathbf{k}, t)$, which are related to the transition amplitude $a_m(\mathbf{k}, t)$. In our numerical calculations, we determine the radiated power of each harmonic frequency by coherently superposing the time derivatives of the intra- and interband currents and then taking the squared modulus of its Fourier transform, i.e., $I_{\text{HHG}}(\omega) = \omega^2 |\text{FT}[\mathbf{J}_{\text{er}}(t) + \mathbf{J}_{\text{ra}}(t)]|^2$.

V. KELDYSH APPROXIMATION AND QUASICLASSICAL ANALYSIS

We may gain more insight into the physics of HHG by applying the so-called Keldysh approximation, as discussed for instance by Vampa *et al.* [67]. This approximation in a crystal solid reads $w(\mathbf{k}, t) = n_v(\mathbf{k}, t) - n_c(\mathbf{k}, t) \approx 1$. Essentially, this means that the population transferred to the conduction band is very small compared to that remaining in the valence band. This approximation is very similar, but never equal, to the one used in the strong field approximation (SFA), developed originally for atoms and molecules [13–15]—we will thus term it SFA in the following. We focus on the discussion of the interband current in this section.

A. Interband current

In this approximation, Eq. (23) decouples from Eq. (24), and we obtain a closed expression for the i th vectorial component ($i = x, y$) for 2D materials of the interband

current,

$$J_{\text{er}}^{(i)}(t) = -i \sum_j \frac{d}{dt} \int_{t_0}^t dt' \int_{\text{BZ}} d^2\mathbf{K} d_{cv}^{(i)}(\mathbf{K} + \mathbf{A}(t)) \times d_{cv}^{(j)}(\mathbf{K} + \mathbf{A}(t')) E^{(j)}(t') \times e^{-iS(\mathbf{K}, t, t') - (t-t')/T_2} + \text{c.c.}, \quad (30)$$

where $S(\mathbf{K}, t, t')$ is the so-called quasiclassical action for the electron-hole, which is defined as

$$S(\mathbf{K}, t, t') = \int_{t'}^t [\varepsilon_g(\mathbf{K} + \mathbf{A}(t'')) + \mathbf{E}(t'') \cdot \tilde{\xi}_g(\mathbf{K} + \mathbf{A}(t''))] dt''. \quad (31)$$

Here, $j = x, y$ indicates the component of the electric field and transition-dipole product which depends on the polarization of the driving laser.

This integral form of the interband current has a nice physical interpretation in terms of the electron-hole (e-h) pair:

(i) At time t' , the e-h pair is excited by the driving laser from the valence band to the conduction band through the dipolar interaction $\mathbf{E}(t') \cdot \mathbf{d}_{cv}(\mathbf{K} + \mathbf{A}(t'))$ at the canonical crystal quasimomentum \mathbf{K} .

(ii) The e-h pair propagates in the conduction band and valence band, respectively, between t' and t , and modifies their trajectories and energies according to Eq. (37) below.

(iii) At time t , the electron has a probability to recombine (or annihilate) with the hole, at which point it emits its excess energy as a high energy-photon.

The expressions Eqs. (30) and (31) are the direct analogues of the Landau-Dykhne formula for HHG in atoms, derived in Ref. [15], following the idea of the simple man's model [14] (see also Ref. [13] for a recent review). Below we will analyze these expressions using the saddle-point approximation over crystal momentum to derive the effects of the Berry curvature on the relevant trajectories.

1. Bloch wave gauge invariance: Quasiclassical action

Before turning to the quasiclassical analysis, let us first demonstrate that the expression for the interband current Eq. (30), together with the action Eq. (31), is gauge invariant under the Bloch wave function transformation (BWT). This refers to the phase freedom of the band eigenstates $|u_{m,\mathbf{k}}\rangle$, whose phases can be chosen independently of each other. As such, a local phase change of the form

$$|u_{m,\mathbf{k}}\rangle \rightarrow |\tilde{u}_{m,\mathbf{k}}\rangle = e^{-i\varphi_m(\mathbf{k})} |u_{m,\mathbf{k}}\rangle \quad (32)$$

should leave the physics unchanged. However, this is not that simple, as multiple relevant quantities change in response to such a BWT. Most notably, the off-diagonal matrix elements $d_{cv}^{(i)}(\mathbf{k}) = |d_{cv}^{(i)}(\mathbf{k})| e^{-i\phi_{cv}^{(i)}(\mathbf{k})}$ ($i = x, y$) of the dipole moment,

$$\tilde{\mathbf{d}}_{cv}(\mathbf{k}) = e^{i\varphi_c(\mathbf{k})} \mathbf{d}_{cv}(\mathbf{k}) e^{-i\varphi_v(\mathbf{k})}, \quad (33)$$

acquire a nontrivial phase shift,

$$\tilde{\phi}_{cv}^{(i)}(\mathbf{k}) = \phi_{cv}^{(i)}(\mathbf{k}) - \varphi_g(\mathbf{k}), \quad (34)$$

for $\varphi_g(\mathbf{k}) = \varphi_c(\mathbf{k}) - \varphi_v(\mathbf{k})$, while the diagonal elements—the Berry connection, $\tilde{\xi}_m(\mathbf{k}) = \mathbf{d}_{mm}(\mathbf{k})$ —shift by the gradient of

the phase function

$$\tilde{\xi}_m(\mathbf{k}) = \xi_m(\mathbf{k}) + \nabla_{\mathbf{k}} \varphi_m(\mathbf{k}). \quad (35)$$

The Berry curvature $\Omega_m(\mathbf{k}) = \nabla_{\mathbf{k}} \times \xi_m(\mathbf{k})$, of course, is not affected.

Since our expressions Eqs. (30) and (31) for the interband current depend explicitly on these gauge-dependent quantities, we are obliged to confirm that the current itself (a physical observable) is not affected by the transformation, as well as to examine the transformation properties of the internal components of those expressions. To do this, we rewrite the expressions above by explicitly taking into account the phases of the transition dipole moment,

$$J_{\text{er}}^{(i)}(t) = -i \sum_j \frac{d}{dt} \int_{t_0}^t dt' \int_{\text{BZ}} d^2\mathbf{K} |d_{cv}^{(i)}(\mathbf{K} + \mathbf{A}(t))| \times |d_{cv}^{(j)}(\mathbf{K} + \mathbf{A}(t'))| E^{(j)}(t') \times e^{-iS(\mathbf{K}, t, t') - (t-t')/T_2 + i(\phi_{cv}^{(j)}(\mathbf{K}, t) - \phi_{cv}^{(i)}(\mathbf{K}, t))} + \text{c.c.}, \quad (36)$$

where the action is “corrected” $\tilde{S}^{(j)}(\mathbf{K}, t, t')$ by the time-dependent dipole phases as

$$\tilde{S}^{(j)}(\mathbf{K}, t, t') = \int_{t'}^t \left[\varepsilon_g(\mathbf{K} + \mathbf{A}(t'')) + \mathbf{E}(t'') \cdot \tilde{\xi}_g(\mathbf{K} + \mathbf{A}(t'')) - \frac{d}{dt''} \phi_{cv}^{(j)}(\mathbf{K} + \mathbf{A}(t'')) \right] dt''. \quad (37)$$

Here $j = x, y$ determines the component of the dipole moment whose phase gets incorporated into the action, determined in part by the polarization of the driving laser. Thus, it is important to note that for nontrivial 2D polarizations this action can split into several (though related) versions, each for a different component of the transition. Note that the dipole phase in the action has the same index as the electric field j . The additional term $\phi_{cv}^{(j)}(t) - \phi_{cv}^{(i)}(t)$ only contributes in the perpendicular-emission configuration $i \neq j$, if one assumes the laser field is linearly polarized along the x direction.

The total action \tilde{S} can be written as

$$\tilde{S}^{(j)}(\mathbf{K}, t, t') = \int_{t'}^t \left[\tilde{\varepsilon}_g(\mathbf{K} + \mathbf{A}(t'')) + \mathbf{E}(t'') \cdot \tilde{\xi}_g(\mathbf{K} + \mathbf{A}(t'')) - \frac{d}{dt''} \tilde{\phi}_{cv}^{(j)}(\mathbf{K} + \mathbf{A}(t'')) \right] dt'', \quad (38)$$

where $\tilde{\varepsilon}_g = \varepsilon_g$ is gauge invariant, but $\tilde{\xi}_g$ and $\tilde{\phi}_{cv}^{(j)}$ are not. However, we can use the transformation rules Eqs. (34) and (35), for the dipole phase and the Berry connection:

$$\tilde{S}^{(j)}(\mathbf{K}, t, t') = \int_{t'}^t \left[\varepsilon_g + \mathbf{E}(t'') \cdot (\xi_g + \nabla_{\mathbf{k}} \varphi_g) - \frac{d}{dt''} (\phi_{cv}^{(j)} - \varphi_g) \right] dt''. \quad (39)$$

(For simplicity, we have dropped the dependence on the variable $\mathbf{K} + \mathbf{A}(t)$ here.) Moreover, by using the attribute that for any function $f(\mathbf{K} + \mathbf{A}(t))$,

$$\frac{d}{dt} f(\mathbf{K} + \mathbf{A}(t)) = -\mathbf{E}(t) \cdot \nabla_{\mathbf{k}} f(\mathbf{K} + \mathbf{A}(t)), \quad (40)$$

as well as the property $\nabla_{\mathbf{k}} = \nabla_{\mathbf{K}}$ [5], we have

$$\left(\mathbf{E}(t'') \cdot \nabla_{\mathbf{K}} + \frac{d}{dt''} \right) \phi_g = 0, \quad (41)$$

which leads finally to the relation

$$\tilde{S}^{(j)}(\mathbf{K}, t, t') = \int_{t'}^t \left[\varepsilon_g + \mathbf{E}(t'') \cdot \boldsymbol{\xi}_g - \frac{d}{dt''} \phi_{cv}^{(j)} \right] dt''.$$

Since this expression is equal to Eq. (37), we find that $\tilde{S}^{(j)}(\mathbf{K}, t, t')$ is thus Bloch-gauge invariant. Both the HHG spectrum and the currents, as physical observables, should be gauge-invariant quantities. Our analysis shows that this is indeed the case in our theoretical model, even in the presence of a nontrivial Berry connection.

In addition, we can also rewrite the semiclassical action as

$$\tilde{S}^{(j)}(\mathbf{K}, t, t') = \int_{t'}^t \left[\varepsilon_g(\mathbf{K} + \mathbf{A}(t'')) + \mathbf{E}(t'') \cdot (\boldsymbol{\xi}_g(\mathbf{K} + \mathbf{A}(t''))) + \nabla_{\mathbf{K}} \phi_{cv}^{(j)}(\mathbf{K} + \mathbf{A}(t'')) \right] dt'', \quad (42)$$

by explicitly calculating the time derivative of the dipole phase, $\frac{d}{dt''} \phi_{cv}^{(j)}$, to bring to the front the explicit factor of the electric field, $\mathbf{E}(t'')$. Here the action phase includes geometric and topological properties, such as the Berry connection and the dipole phase $\phi_{vc}^{(j)}(\mathbf{k})$. Given this form, it is probable that the electron-hole pair trajectories will be modified by these geometric properties, thereby affecting the non-linear HHG process. The total phase of Eq. (36) has an extra dipole-phase contribution, $\phi_{cv}^{(i)} - \phi_{cv}^{(j)}$ regarding the ‘‘perpendicular’’ interband current to the electric field. From the phase transformation given by Eq. (34), this phase difference term is also gauge invariant.

B. Quasiclassical approach and electron-hole pair trajectories

Assuming that the exponentiated quasiclassical action $e^{-iS} = e^{-iS(\mathbf{K}, t, t')}$ oscillates rapidly as a function of the crystal momentum \mathbf{K} , one can apply the saddle-point approximation to find the points \mathbf{K}_s where the integrand’s contributions to the interband current Eq. (36) concentrate. These are solutions of the saddle-point equation $\nabla_{\mathbf{K}} S(\mathbf{K}, t, t')|_{\mathbf{K}_s} = \mathbf{0}$, which can be rephrased as

$$\Delta \mathbf{x}_c(\mathbf{K}_s, t, t') - \Delta \mathbf{x}_v(\mathbf{K}_s, t, t') = \mathbf{0}. \quad (43)$$

From the last equation, two different trajectories are identified, the first one related to the excited electron $\Delta \mathbf{x}_c(\mathbf{K}_s, t, t')$ in the conduction band and the second one regarding the trajectory $\Delta \mathbf{x}_v(\mathbf{K}_s, t, t')$, followed by the hole in the valence band. We then obtain a general m th trajectory for the electron ($m = c$) and hole ($m = v$), which reads

$$\Delta \mathbf{x}_m(\mathbf{K}_s, t, t') = \int_{t'}^t \left[\mathbf{v}_{gr,m} + \mathbf{E}(t'') \times \boldsymbol{\Omega}_m + (\mathbf{E}(t'') \cdot \nabla_{\mathbf{K}}) \left(\boldsymbol{\xi}_m + (-1)^m \frac{1}{2} \nabla_{\mathbf{K}} \phi_{cv}^{(j)} \right) \right] dt'', \quad (44)$$

where $(-1)^m$ is the alternating sign $(-1)^c = +1$ and $(-1)^v = -1$, and the group velocity of the m th band is $\mathbf{v}_{gr,m} = \nabla_{\mathbf{K}} \varepsilon_m$. Here we recognize the Berry curvature $\boldsymbol{\Omega}_m$ as well as the

anomalous velocity $\mathbf{v}_{a,m}$, which are given by $\boldsymbol{\Omega}_m = \nabla_{\mathbf{K}} \times \boldsymbol{\xi}_m$ and $\mathbf{v}_{a,m} = \mathbf{E}(t) \times \boldsymbol{\Omega}_m$, respectively, for the electron-hole trajectories of Eq. (44). We can rewrite the previous expression as

$$\begin{aligned} \Delta \mathbf{x}_m(\mathbf{K}_s, t, t') &= \int_{t'}^t \left[\mathbf{v}_{gr,m} + \mathbf{v}_{a,m} - \frac{d}{dt''} \left(\boldsymbol{\xi}_m + \frac{(-1)^m}{2} \nabla_{\mathbf{K}} \phi_{cv}^{(j)} \right) \right] dt''. \end{aligned} \quad (45)$$

These electron-hole pair trajectories, together with the saddle-point condition of Eq. (43), should produce complex-valued solutions for \mathbf{K}_s , as is the case for HHG of gases.

However, finding the solutions \mathbf{K}_s is not a trivial task, since it depends explicitly on the geometrical features, i.e., the Berry curvature and connection, and the phase of the dipole matrix elements. Moreover, this gets further complicated as the eigenstates and eigenvalues that make up the energy bands are expected to exhibit branch cuts and branch points connecting the two bands [68,69]. Once the momentum is allowed to take on complex values (as it does in complex band structure theory [70]), this leads to a nontrivial geometrical problem with a high dimensionality whose analysis requires detailed attention.

Nevertheless, these saddle points \mathbf{K}_s should have a component perpendicular to the driving laser field $\mathbf{E}(t)$ (for the case of linear drivers), which appears as a consequence of anomalous-velocity features and in particular of the Berry curvature $\boldsymbol{\Omega}_m(\mathbf{k})$.

VI. HHG IN PRACTICE: THE ROLE OF INTER- AND INTRABAND CURRENTS

Having developed the theory of HHG in topological solids over the previous sections, we now turn to applying it to some examples of topologically trivial and nontrivial systems. In a semiconductor or insulator driven by midinfrared lasers or THz sources, the harmonic emission is governed by the coherent sum of the intraband $\mathbf{J}_{ra}(t)$ and interband $\mathbf{J}_{er}(t)$ current oscillations $\mathbf{J}(t) = \mathbf{J}_{ra}(t) + \mathbf{J}_{er}(t)$ [18,19] given above by Eqs. (27) and (29). We stress, as mentioned previously, that the interband current contains explicit information about the Berry curvature through the cross product of the dipole moments. This is important, since in 2D materials the integral of the Berry curvature over the BZ is the topological invariant, i.e., the Chern number. In Fig. 1(Y), we depict a cartoon of the HHG physical process which takes place in topological materials.

Via our quasiclassical saddle-point analysis we find the following: (1) The electron-hole pair is more likely to be excited around the K' point than the Γ or K points, since the energy gap is larger at those points compared to K' for this HM. (2) The electron-hole pair is propagated by the driving laser field along the path indicated with black dashed line of the BZ emitting intraband harmonics in the process. This propagation follows the laser vector potential in the momentum plane, with the real-space motion (and thus the intraband current) governed by the group velocity produced by the energy-band dispersion as well as an anomalous velocity caused by the Berry curvature. (3) Finally, the electron can recombine with




FIG. 3. (a) Experimental [27] and (b) theoretical HHG spectra from an α -quartz crystal. Both the theoretical and experimental data are averaged along the parallel and perpendicular harmonic emission with respect to a MIR 800 nm laser linearly polarized along the Γ - K crystal orientation [cf. Fig. 1(b)]. (c) Parallel and (d) perpendicular harmonic emission from our α -quartz toy model, showing the interband (blue), intraband (red), and total (green) currents. We use laser parameters following Ref. [27], with a wavelength $\lambda_0 = 800$ nm giving a central frequency $\omega_0 = 0.056$ a.u. (1.53 eV), FWHM duration of ten cycles under a Gaussian envelope, and peak intensity of $I_0 = 3.2 \times 10^{12}$ W/cm². We use a honeycomb lattice with lattice constant $a_0 \sim 4.8$ Å, NN hopping $t_1 = 2.1$ eV, NNN hopping amplitude $t_2 = 0.36$ eV with magnetic flux $\phi_0 = 0$, and staggering potential $M = 4.5$ eV, with a band gap $E_g \sim 9$ eV, and a dephasing time of $T_2 = 220$ a.u. = 5.3 fs.

its hole and release its energy as a photon. This interband emission carries traces of the topological features of the material via Berry-curvature contributions to the action, which are accrued in the continuum traversal as well as via the dipole matrix elements that mediate the transition.

We start by comparing our theory to the results of the recent HHG experiment of Luu and Wörner [27] on α -quartz (SiO₂). Since a cut along the z direction of this crystal has an approximate honeycomb lattice structure, we use the trivial limit of the HM to validate our theory. Due to the breaking of IS in α -quartz, the HHG spectrum contains even HOs, shown in Fig. 3(a), mainly along the perpendicular emission configuration. As we detail below, our simulations show that the interband contribution has a larger magnitude than the intraband current. Generally, the experimental features of Ref. [27] are qualitatively reproduced by our theory [see Fig. 3(b)]; moreover, this approach qualitatively agrees with Liu *et al.*'s experiment reported for MoS₂ [26]. That said, it is important to note that the calculations overestimate the intensity yield for the HOs of the plateau.

The discrepancy between theory and experiment is related to the limitations of this toy model under the tight-binding approach, which will not recreate the whole experimental complexity for the following reasons: (i) the lattice crystal is not a monolayer (the crystal thickness is $20 \mu\text{m}$ [27]), (ii) the crystalline structure of the hexagonal 3D lattice of SiO₂ is much more complex than our simplified 2D lattice, (iii) the possibility of higher conduction bands and lower valence bands playing a role is not considered in our

approximations [71], and (iv) the propagation effects in the $20\text{-}\mu\text{m}$ -thick crystal can play a fundamental role [72]. However, despite this discrepancy, we believe that physical insight can be extracted in terms of identifying dominant mechanisms for the harmonic emission (intraband)—nonlinear Bloch oscillations of charge carriers [24], and (interband) electron-hole recombination processes [19]—and their relative importance.

We now turn to the parallel and perpendicular components of the harmonic emission when driven by a linearly polarized laser, with respect to the driver's polarization, which we show in Figs. 3(c) and 3(d) broken out into the intra- and interband components. First, our calculations show the same qualitative features of the experiments [26,27]: odd and even harmonics are emitted along the parallel and perpendicular emission directions, respectively. Second, the interband current dominates over the intraband one for both parallel and perpendicular configurations, at least for these excitation conditions, and with the usual caveats regarding the nontrivial coupling between the two components [24]. This implies that a recollision mechanism is likely to be at play in a system where the IS is broken. We also note that along the perpendicular emission, HHG shows only even harmonics for the intraband analysis exhibiting dominant behavior for largest HOs. We believe that the appearance of even harmonics along the perpendicular direction is not only a matter of intraband or Berry-curvature effects: they are produced by the interband process, due to a combination of the dipole phases, Berry curvature, and Berry connection features. This explanation is apparent both in our saddle-point arguments for the interband




FIG. 4. High-order harmonics for trivial and nontrivial topological materials. We show the harmonic emission from equal band-gap instances from the trivial and topological phases driven by lasers with (a) linear polarization along Γ - K and (b) right-circular polarization; the violet areas indicate the difference in the emission between the two phases. We use $t_1 = 2.04$ eV, $t_2 = t_1/3$ and $M_0 = 2.54t_2$, with lattice constant $a_0 = 1$ Å, driven by a laser centered at 3.25 μm ($\omega_0 = 0.014$ a.u.) with a Gaussian envelope with 14-cycle FWHM and a peak electric-field strength of $E_0 = 0.0045$ a.u. ($I_0 = 7.0 \times 10^{11}$ W/cm 2), and we use a dephasing time of $T_2 = 5.2$ fs. The magnetic flux of the trivial and nontrivial phases, $\phi_0 = 0.06$ and $\phi_0 = 1.16$ rad, respectively, are chosen to give equal band gaps of $E_g \approx 3.0$ eV.

mechanism above and the numerical calculations given here. The HHG emission is not only the product of a single effective intraband contribution; both interband and intraband mechanisms are present in the rich physics of this nonlinear harmonic emission process.

Finally, as additional validation of our simulations, we have studied the scaling of the harmonic cutoff with respect to the driving-field amplitude, which we report below in Appendix B. As expected, the cutoff exhibits a linear scaling in good agreement with previously obtained theoretical and experimental results [18,22].

VII. SIGNATURES OF TOPOLOGICAL PHASES IN HHG

We now address the question of whether HHG is sensitive to (i) different topological phases and (ii) topological transitions. To this end, in Fig. 4 we show the calculated full HHG radiation spectrum, $I_{\text{HHG}}(\omega) = \omega^2(|J_x(\omega)|^2 + |J_y(\omega)|^2)$ produced by linearly polarized [Fig. 4(a)] and circularly polarized [Fig. 4(b)] driving lasers, comparing instances from the trivial and the topologically nontrivial phase at equal band gaps. For a linearly polarized driving laser, the harmonic intensity spectrum exhibits relatively small differences between the two phases (though signatures of the topological phase do appear in the polarization state of the harmonics [44]). In our results, we observe an enhancement of about one order of magnitude for even-order harmonics in the low spectral region, i.e., the second, fourth, and sixth harmonics.

For a circularly polarized driver, on the other hand, shown in Fig. 4(b), the spectra produced by the different topological phases exhibit far greater differences, in particular: (1) a considerable enhancement about four orders of magnitude in plateau harmonics in the topological phase as compared with the trivial phase, (2) a shorter cutoff for the topological emissions in comparison to the trivial, and (3) an asymmetric yield between the corotating ($3n + 1$) and contrarotating ($3n + 2$) HOs in plateau harmonics. As such, it is clear that circularly polarized drivers have significant potential in bringing out the signatures of the topological phase in the harmonic spectrum and we will focus on them for the rest of this work.

The circularly driven spectra of Fig. 4(b) are also noticeably different from the HHG spectra driven by linearly polarized pulses, in that they exhibit a clear selection rule, where harmonics of order $3n$ are forbidden and $3n \pm 1$ are allowed, for an integer n . This selection rule is identical to that observed in the well-known bicircular field configuration—two counter-rotating circularly-polarized drivers at frequencies ω_0 and $2\omega_0$ [73]—and it stems from the same origin, a dynamical symmetry of the system, which combines a rotation by 120° with a time delay [74]. This symmetry argument has been applied in solids [55,75–78] but has also been applied to molecules [79–81] and even plasmonic metamaterials [82].

In our case, however, the generating medium—the HM on a hexagonal lattice—is symmetric under 120° rotations, but it lacks reflection symmetry, which allows it to respond nontrivially to the different helicities of circularly polarized driving lasers. This is much like p states in noble-gas atoms [83,84] and chiral molecules [85,86], which points to the CD of the optical response (i.e., the difference in response to left- and right-handed circularly polarized drivers) as a natural place to look for signatures of the topological phase of the material.

VIII. CIRCULAR DICHROISM

As an initial test of this CD idea, we show in Figs. 5(a) and 5(b) the total harmonic spectra produced in topologically trivial and nontrivial phases, respectively, when driven by RCP and LCP (shown in blue and red, respectively) laser pulses. The topologically nontrivial phase shows a distinct increase in HO response of the form $k = 3n + 1$ (which corotates with the driving laser due to the usual selection rules) when driven by an RCP laser as compared with an LCP driver, while the topologically trivial phase shows a small *decrease* in the emission of those harmonics. (In absolute terms, this combination of laser helicities matches that of the next-nearest-neighbor coupling $t_2 e^{i\phi_0}$, as shown in Fig. 1(c); if a negative ϕ_0 is chosen instead, the effect reverses direction.)

This observation indicates that, indeed, the CD of the harmonic emission carries clear information about the topological phase performing the emission (the CD is invariant under the electromagnetic minimum-coupling gauge), and we




FIG. 5. Circular dichroism as a signature of topological phase transitions in HHG. (a), (b) Harmonic spectra of equal band-gap instances from the trivial and the topological phase, respectively, driven by right- and left-circularly polarized lasers, with the difference between the two emissions shaded in green. (We use the same parameters as in Fig. 3.) (c), (d) Circular dichroism (CD), as defined in Eq. (46), for the magnetic-flux and staggering potential parameter scans marked with dashed lines in Fig. 1, for the plateau harmonics. The top horizontal axes show the energy band gap E_g in atomic units (a.u.) as a function of ϕ_0 (c) and M_0/t_2 (d), respectively. The trivial and topological regions are shaded green and blue, with the critical region shaded red.

focus on it for the rest of this paper. We define, in particular,

$$\text{CD}_k = \frac{I_{k,\text{RCP}}^{(+)} - I_{k,\text{LCP}}^{(-)}}{I_{k,\text{RCP}}^{(+)} + I_{k,\text{LCP}}^{(-)}}, \quad (46)$$

that is, the normalized difference in harmonic response driven by left- and right-handed circularly polarized drivers, for HOs $k = 3n + 1$. (For the counter-rotating harmonics $k = 3n + 2$, the signs in the superscripts should be reversed.) Here the CD is defined in terms of the circular components of the harmonic emission, $I_{k,v}^{(\pm)} = (k\omega_0)^2 |J_{x,v}(k\omega_0) \pm iJ_{y,v}(k\omega_0)|^2$, $J_{x,v}$ and $J_{y,v}$ denoting the x and y components of the spectral current driven by a $v = \text{RCP, LCP}$ laser.

Our main result is shown in Figs. 5(c) and 5(d). These plot the CD for the corotating plateau harmonics over the three distinct topological phases of the Haldane phase diagram [see Fig. 1(a)]. (i) in Fig. 5(c), we scan over ϕ_0 with M_0 constant [see horizontal cut depicted in orange dashed line of Fig. 1(a)]. And (ii) in Fig. 5(d) at $\phi_0 = \pi/2$, we also scan the CD over M_0/t_2 [vertical cut depicted in blue dashed line of Fig. 1(a)]. For case (i), we observe two clear topological-plateau structures in the circular dichroism with values of $\text{CD} \approx \mp 1$ for the topological invariants $C = \mp 1$, a clear *topological-plateau signature for the CD of the corotating harmonics*. Particularly, high oscillatory behavior is observed in the CD for the trivial phase $C = 0$, which deviates from the robust CD topological-plateau structure, once the topological phase boundary is

crossed. In short, the CD observable shows clear distinct characteristics as a function of the Chern number $C = -1, 0, +1$.

In case (ii), while the $\text{CD} \approx +1$ for $C = +1$, once the topological phase boundary is crossed, the CD disappears relatively quickly as M_0/t_2 increases. This effect can be understood rather simply, since in the $M_0 \gg t_2$ limit, the material's Hamiltonian is dominated by the staggering potential M_0 , which is reflection symmetric and does not support a CD response. However, the most notable aspect of the CD spectra is that it still exhibits clear variations depending on the topological phase the system is in. This is true even for instances of the HM which are equally chiral at the level of the Hamiltonian (in the sense of having identical M_0/t_2 ratios). In other words, the high-harmonic CD response shows a clear signature of the topological phase of the system.

There is also a clear transition regime between the two topological phases, shown shaded pink in Figs. 5(c) and 5(d), where the CD (particularly in the M_0/t_2 scan) takes on irregular values. At the topological phase boundary the band gap closes, as it must do whenever the Chern number changes value, and at these points the material becomes a semimetal with one exact graphene-like Dirac cone per BZ. In the critical region around those points [see Fig. 1(Q)], the material is a semiconductor with a negligible gap, and it is similarly permissive to nonadiabatic transitions. Within this critical region, as in graphene [77], the lowered band gap produces a much higher e-h population, which naturally raises the harmonic emission. This is shown in Fig. 6, which displays the total




FIG. 6. Variation of the harmonic spectrum for circularly-polarized drivers. (a), (b) Harmonic spectrum for the magnetic-flux parameter scan indicated with the red dashed line in Fig. 1(a), driven by LCP and RCP light, respectively, with ϕ_0 ranging over a 60-point scan of the interval $[0, \pi]$ and the staggering potential held fixed at $M_0 = 2.54t_2$. At the critical region, where the band gap closes between the two phases [as indicated in (c), which charts the band gap], the harmonic emission increases, as carrier creation is much easier. The signature of the topological phase is in the circular dichroism [the difference between the RCP-driven signal in (b) and the LCP-driven signal in (a)], which we plot in Fig. 5 for the harmonics in the green box. (d)–(e) Identical plots for the staggering-potential scan marked by the blue dashed line in Fig. 1(a), varying M_0/t_2 at fixed magnetic flux of $\phi_0 = \frac{\pi}{2}$. The (c) and (f) depict the band gaps, E_g , as a function of ϕ_0 and M_0/t_2 , respectively.

emission for both of the phase-space scans of Fig. 5 for each of the driving helicities.

This enhanced emission acts naturally as a litmus test that the topological phase boundary itself has been reached, to complement the CD as a signature of which topological phase the material is in. Moreover, the total-emission scans of Fig. 6 confirm that the features observed previously also hold more broadly over the phase diagram: for one, the topologically nontrivial phase shows a higher emissivity than the trivial material, at similar band gaps [shown in Figs. 6(c) and 6(f)], and, similarly, there are noticeable changes in the harmonic cutoff as well.

IX. DEPHASING TIME IN THE HARMONIC EMISSION FOR TOPOLOGICAL PHASES

It is also important to point out that the features previously shown both in the harmonic intensity and the CD depend (sometimes sensitively) on the dephasing time T_2 in our calculation; we now turn to examine this dependence. Since the electron-hole trajectory mechanism is modified by the dephasing time, particularly for long trajectories, the harmonic intensity, and from it the CD, inherit this effect, as expected [19,22,67]. On the other hand, and somewhat surprisingly, in the topological phase the CD is more robust against

dephasing than in the trivial phase, particularly for the $k = 3n + 1$ corotating harmonics, which again points to a clear role of the material's topology in its harmonic response, and cements the interaction between the two as a clear target for continued investigation.

We now study the role of the dephasing time T_2 on the CD for the trivial and topological phases. Figures 7(a) and 7(b) show the HHG spectra as a function of T_2 for trivial and nontrivial topological phases, respectively. The emitted harmonic intensity is sensitive to the dephasing time, as shown also in literature for linearly polarized lasers [67]. Here we also observe sensitivity of the HHG emission yields with respect to T_2 for both trivial and topological phases. T_2 usually reduces the coherence of the occupations and therefore also the electron-hole recombination mechanisms, more strongly for long trajectories [67].

These effects also appear to take place in topological phases under this simple two-band approximation, which would lead to a potential dependence of the dichroism for the different HOs. Additionally, we computed the HHG spectra for the case $T_2 \rightarrow \infty$ (not shown, for simplicity), in which case the harmonic plateau becomes extremely noisy, making it impossible to extract any reliable insights from those simulations. Similar features can be also noticed in the numerical background noise of Fig. 7(b) as the dephasing time increases.




FIG. 7. Role of the dephasing time in the high-harmonic emission and in its circular dichroism. (a), (b) High-harmonic generation spectra for various dephasing times T_2 , for trivial and topological phases, respectively, driven by a right-circularly polarized laser. (c)–(f) Circular dichroisms for $3n + 1$ (corotating) and $3n + 2$ (contrarotating) harmonic orders [(c)–(f), respectively], for trivial (c), (e) and topological (d), (f) phases. The parameters for the material and the driving laser are the same as in Fig. 2 with a slightly altered peak field strength, $E_0 = 0.005$ a.u., and pulse duration, $N_c = 16$.

For the trivial phase, Figs. 7(c) and 7(e) show the dichroism as a function of T_2 for $3n + 1$ (corotating) and $3n + 2$ (counter-rotating) HOs, respectively; we find a strong dependence of the CD on the dephasing time for each individual order. Nevertheless, when averaging the CD for corotating orders over T_2 , it remains negative, providing a systematic tendency around the plateau and, likely, the cutoff. In contrast, for the counter-rotating orders, CD oscillates between positive and negative values depending on lowest or highest region of the spectrum. This behavior seems to be a natural consequence of the complex relationship between the electron-hole recombination mechanism, intraband and interband mechanisms, and the T_2 .

In contrast, the corotating CD for the topological phase shown in Fig. 7(d) is much more robust against T_2 , exhibiting values around +1, most noticeably when $\text{HO} > 4^{\text{th}}$. For counter-rotating orders [see Fig. 7(f)], the CD describes similar features across the whole spectrum as in the trivial phase. In that sense, the CD depends on the region of the harmonic spectrum at play. These observations show that the CD is a more robust quantity for the corotating than for the counter-

rotating HOs, which is our main reason for focusing on the CD of the former in the previous section.

X. CONCLUSIONS AND OUTLOOK

In summary, our theory shows that the HHG spectrum is (i) able to probe topological phase transitions, (ii) able to test topological invariants by using the CD of the corotating harmonics, and (iii) extremely sensitive to the breaking of both time-reversal and inversion symmetries (TRS, IS, respectively).

Our results are applicable to the HM, so they constitute clear evidence that the HHG process can trace topological phases and transitions in a topological Chern insulator. But it is still necessary to explore other classes of Chern insulators as well as topological insulators to confirm that this HHG-circular dichroism observable is a broadly defined quantity to map topological transitions. In particular, we show CD calculations for the so-called Wilson mass model [87] (WMM), which suggests that the results obtained by the HM are generally extended to Chern insulators. Since the main focus of

this paper is on HM, we will show partially the regarding evidences for WMM in Appendix A. The formulation of relativistic fermions in lattice gauge theories [88] is hampered by the fundamental problem of species doubling [89,90], namely, the rise of spurious fermions that modify the physics at long wavelengths. To prevent the abundance of such fermion doublings, a suitable tailoring of their masses is required, leading to the so-called Wilson fermions [87]. The WMM with an inverted mass gives rise to a certain axion background [91–97], and thus may help explain the lack of experiments confirming the charge-parity violation in strong interactions (for quantum simulations of this model, see Ref. [98]). When exposed to ultraintense short laser pulses, the WMM shows similar behavior to the one we report here for the HM; these results will be reported in a future publication. In that general sense, our theory can be applied to a larger range of topological materials with similar band-gap properties described in this paper, as well as extended to THz sources [99]. Another possible application could be Bi_2Se_3 , which is a good candidate for topological insulators; possible layer-thick modifications could be created in this material to control the topological transition as in Ref. [100]. Also, THz sources should open a path to access HHG in topological materials by pushing the driving photon energy well below the band gap in two different topological orders; this would also raise new questions about exploring the dynamics in strongly correlated systems with spin-orbit couplings and degeneracies, in particular in bilayer systems and twistrionics [37,101–104].

Finally, it is important to note that all of these ideas can be applied and tested in quantum simulators [105], in particular, using ultracold atoms and lattice shaking [106], as well as Rydberg atoms, polaritons, or circuit QED [107–112]. The use of quantum simulators to study strong-field phenomena is an emerging application of the former [113–115], and it is naturally suited to the study of systems in ordered lattices, so it should provide additional clarity to the role of material topology in high-harmonic emission.

The data and code used in this paper have been deposited in the GitHub public repository [116].

ACKNOWLEDGMENTS

We thank T.T. Luu and H.J. Wörner for providing experimental data in an editable format. A.C. thanks D. Baykusheva for running extra calculations of Figs. 5 and 6, in the supercomputer Sherlock at SLAC, and for useful discussions of the CD results, also to N. Aguirre and A. Adedoyin for suggestions in technical details of numerical implementations. A.D., E.P., A.S.M., A.P., M.F.C., and M.L. acknowledge support from ERC AdG NOQIA, Spanish Ministry of Economy and Competitiveness (“Severo Ochoa” program for Centres of Excellence in R&D (CEX2019-000910-S), Plan National FISICATEAMO and FIDEUA PID2019-106901GBI00/10.13039/501100011033, FPI), Fundació Privada Cellex, Fundació Mir-Puig, and from Generalitat de Catalunya (AGAUR Grant No. 2017 SGR 1341, CERCA program, QuantumCAT_U16-011424, co-funded by ERDF Operational Program of Catalonia 2014-2020), MINECO-EU QUANTERA MAQS (funded by State Research Agency (AEI) PCI2019-111828-

2/10.13039/501100011033), EU Horizon 2020 FET-OPEN OPTOLogic (Grant No. 899794), and the National Science Centre, Poland-Symfonia Grant No. 2016/20/W/ST4/00314. E.P. acknowledges CELLEX-ICFO-MPQ fellowship funding. A.P. acknowledges funding from the grant ref. RTI2018-097355-A-I00 (MCIU/AEI/FEDER, UE) and TALENTO grant ref. 2017-T1/IND-5432 (Comunidad de Madrid). A.D. acknowledges support of Juan de la Cierva fellowship (IJCI-2017-33180) and the fellowship from “la Caixa” Foundation (ID 100010434) and from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 847648. The fellowship code is (LCF/BQ/PR20/11770012). M.F.C. is thankful for the support of project Advanced Research using High Intensity Laser Produced Photons and Particles (No. CZ.02.1.01/0.0/0.0/16019/0000789) from the European Regional Development Fund (ADONIS) and the Grantová agentura České Republiky (GACR Grant No. 20-24805J). A.S.M. acknowledges funding from the UK Engineering and Physical Sciences Research Council (EPSRC) InQuBATE Grant No. EP/P510270/1. A.C., W.Z., S.K., C.T., and A.S. acknowledge support from LDRD No. 2160587 and the Science Campaigns, Advanced Simulation, and Computing and LANL, which is operated by LANS, LLC for the NNSA of the US DOE under Contract No. DE-AC52-06NA25396 CNLS-LANL. A.C., D.K., and D.E.K. acknowledge support from Max Planck POSTECH/KOREA Research Initiative Program (Grant No 2016K1A4A4A01922028) through the National Research Foundation of Korea (NRF) funded by Ministry of Science, ICT & Future Planning, partly by a Korea Institute for Advancement of Technology (KIAT) grant funded by the Korea Government (MOTIE) (No. P0008763, The Competency Development Program for Industry Specialist). A.C. thanks Alma Dreyer and Joshua Dreyer-Chacón for all their support and their understanding in the extra-time and efforts I needed to finalize this project.

APPENDIX A: HAMILTONIAN FOR THE WILSON MASS MODEL

In the main text, we focused on the HM with IS breaking. One question is, whether or not the observed CD produced by HHG is model dependent. Here, we fully answer that question by introducing the simplest square lattice with two localized orbitals per atomic site (\mathcal{A} and \mathcal{B}) (WMM). The Hamiltonian has IS $\mathcal{P}H_0(\mathbf{k})\mathcal{P}^\dagger = H_0(-\mathbf{k})$, in which \mathcal{P} defines the IS operator for $\mathbf{k} \rightarrow -\mathbf{k}$. Note, nevertheless, this WMM breaks the TRS as well as HM does. The WMM Hamiltonian reads

$$H_0(\mathbf{k}) = \Delta \sin(k_x a_0) \sigma_1 + \Delta \sin(k_y a_0) \sigma_2 - \{2t_1[\cos(a_0 k_x) + \cos(a_0 k_y)] + \mu\} \sigma_3, \quad (\text{A1})$$

where Δ is the hopping parameter between \mathcal{A} and \mathcal{B} . This denotes the overlap integral of the two NNs with localized states \mathcal{A} and \mathcal{B} , which in principle can be purely imaginary. Here μ is the relative on-site potential between states \mathcal{A} and \mathcal{B} of the square Wilson lattice. The lattice constant is a_0 and t_1 is the hopping parameter of the same two-consecutive-localized states $\mathcal{A} - \mathcal{A}$ or $\mathcal{B} - \mathcal{B}$ at the square lattice. We define the so-called Wilson mass \mathcal{M} term [87,94] by $\mu = 2t_1\mathcal{M}$ and




FIG. 8. High harmonics and circular dichroism from Wilson Mass Model for trivial, topological phases and transitions. (a) Total high-harmonic generation emissions for trivial (blue) and topological (red with violet shadow) phases. The Wilson mass model (WMM) parameters of lattice constant $a_0 = 3.33$ a.u. and others for trivial, $C = 0$ (topological, $C = +1$) phase are hopping parameter $t_1 = 0.021$ a.u. (0.034 a.u.) with Wilson mass term $\mathcal{M} = \frac{10}{3}$, ($\frac{6}{5}$) and hopping $\delta_0 = 0.0434$ (0.0723). Those WMM parameters produce equal band gaps of $E_g \approx 3.0$ eV for trivial and the topological phases. (b) Depicts the topological phase transition for corotating harmonic orders ($3n + 1$) by the normalized circular dichroism (CD) as a function of \mathcal{M} at a fixed hopping parameter of $t_1 = 0.034$ a.u. The top horizontal axis points to the energy gap E_g (atomic units are used) as a function of \mathcal{M} . The CD shows two different plateaus of $CD \approx +1$ and $CD \approx -1$ for, respectively, the topological invariants $C = +1$ and $C = -1$. On the contrary, for regions where the Chern number $C = 0$, the CD is oscillating. The latter observations are absolutely consistent with the results provided by Haldane model in Fig. 5(c). The laser parameters are the same as in Fig. 5.

rescale $\Delta = 2t_1\delta_0$ by δ_0 . Here, \mathcal{M} and δ_0 are adimensional parameters which control the topological phases with $C = \pm 1$ and $C = 0$. The calculated high-harmonic spectrum for left and right circularly polarized laser-lights are shown in Fig. 8(a) for two different topological phases, trivial $C = 0$ and topological $C = +1$, respectively. We observe a predominant behavior of the harmonic nonlinear responses from the topological phase, a clear enhancement (two or three orders of magnitude across the harmonic-plateau) and a harmonic cutoff shift too. Finally, we report in Fig. 8(b) calculations of the CD as a function of the mass \mathcal{M} in which three distinct topological phases exist, as well as phase transitions. The CD has values of $CD \approx \pm 1$ for the topological invariants $C = \pm 1$, and CD oscillates for the trivial phase. The latter observations denote three different topological structures for the CD in the three corresponding topological phases. These CD results produced by HHG in the Wilson mass model are clearly consistent with the evidences obtained from HM. Hence, we can conclude that the circular dichroism is a general tool which can map topological phases and transitions in Chern insulators.

APPENDIX B: LINEAR FIELD SCALING OF THE HIGH HARMONIC CUT-OFF

To validate our theoretical model and numerical variable-gauge methods, we investigate how the HHG emission cutoffs

for the total intraband and interband contributions behave as a function of the laser field strengths, for both trivial and topological phases.

Figures 9(a) and 9(b) show the results for the total harmonic emission $I_{\text{HHG}}(\omega) = \omega^2[|J^{(x)}(\omega)|^2 + |J^{(y)}(\omega)|^2]$ of the trivial and nontrivial topological phases. First, we see a clear linear cutoff in terms of the laser strength for the topological trivial case. This tendency is in very good agreement with the pioneering experimental observation of Ref. [18] and the theoretical confirmation by Vampa *et al.* in Refs. [22,67]. In the case of the topological phase, we observe a linear cutoff scaling as a function of the laser field strength, similar to the trivial one.

We also depict, in Figs. 9(c) and 9(d), the corresponding intraband harmonic emissions for trivial and nontrivial phases, via the group and anomalous velocity emissions. We find that the intraband case for the topological emission seems to exhibit a much larger intensity yield than the trivial phase (though the coupling between the two also needs to be considered in more detail [24]).

In contrast, Figs. 9(e) and 9(f) show that, for our excitation conditions, the interband contributions for the trivial and topological phases dominate around the plateau and cutoff in comparison to intraband emissions. One therefore needs to carefully consider and analyze the interband mechanism for high nonlinear emission, i.e., $\text{HO} > 5^{\text{th}}$, and not uniquely the intraband mechanism.




FIG. 9. Trivial and topological high harmonic cutoff law for linear polarized driving fields. Total HHG emissions for trivial and topological phases, with the same HM parameters than those used in Fig. 2, as a function of the laser field strength E_0 are depicted in (a) and (b), respectively. (c), (e) The decomposition of the total harmonic emission for the intraband and interband emissions of the trivial phase, respectively. (d), (f) The intraband and interband current emissions for the topological phase. Green lines mimic the linear cutoff for both trivial and topological phases. The laser field for these HHG calculations is linearly polarized along the $K' - \Gamma - K$ direction of the honeycomb hexagonal lattice.

-
- [1] J. M. Kosterlitz, Nobel lecture: Topological defects and phase transitions, *Rev. Mod. Phys.* **89**, 040501 (2017).
- [2] K. von Klitzing, The quantized Hall effect, *Rev. Mod. Phys.* **58**, 519 (1986).
- [3] H. L. Stormer, Nobel lecture: The fractional quantum Hall effect, *Rev. Mod. Phys.* **71**, 875 (1999).
- [4] D. Haddad, F. Seifert, L. S. Chao, S. Li, D. B. Newell, J. R. Pratt, C. Williams, and S. Schlamminger, Invited article: A precise instrument to determine the Planck constant, and the future kilogram, *Rev. Sci. Instrum.* **87**, 061301 (2016).
- [5] X.-L. Qi and S.-C. Zhang, Topological insulators and superconductors, *Rev. Mod. Phys.* **83**, 1057 (2011).
- [6] M. Z. Hasan and C. L. Kane, Colloquium: Topological insulators, *Rev. Mod. Phys.* **82**, 3045 (2010).
- [7] N. Goldman, G. Juzeliunas, P. Öhberg, and I. B. Spielman, Light-induced gauge fields for ultracold atoms, *Rep. Prog. Phys.* **77**, 126401 (2014).
- [8] T. Ozawa, H. M. Price, A. Amo, N. Goldman, M. Hafezi, L. Lu, M. C. Rechtsman, D. Schuster, J. Simon, O. Zilberberg, and I. Carusotto, Topological photonics, *Rev. Mod. Phys.* **91**, 015006 (2019).
- [9] S. D. Huber, Topological mechanics, *Nat. Phys.* **12**, 621 (2016).
- [10] W. J. M. Kort-Kamp, Topological Phase Transitions in the Photonic Spin Hall Effect, *Phys. Rev. Lett.* **119**, 147401 (2017).
- [11] D. T. Tran, A. Dauphin, A. G. Grushin, P. Zoller, and N. Goldman, Probing topology by “heating”: Quantized circular dichroism in ultracold atoms, *Sci. Adv.* **3**, e1701207 (2017).

- [12] J. Itatani, J. Levesque, D. Zeidler, H. Niikura, H. Pépin, J.-C. Kieffer, P. B. Corkum, and D. M. Villeneuve, Tomographic imaging of molecular orbitals, *Nature* **432**, 867 (2004).
- [13] K. Amini, J. Biegert, F. Calegari, A. Chacón, M. F. Ciappina, A. Dauphin, D. K. Efimov, C. F. de Morisson Faria, K. Giergiel, P. Gniewek *et al.*, Symphony on strong field approximation, *Rep. Prog. Phys.* **82**, 116001 (2019).
- [14] P. B. Corkum, Plasma Perspective on Strong Field Multiphoton Ionization, *Phys. Rev. Lett.* **71**, 1994 (1993).
- [15] M. Lewenstein, P. Balcou, M. Y. Ivanov, A. LHuillier, and P. B. Corkum, Theory of high-harmonic generation by low-frequency laser fields, *Phys. Rev. A* **49**, 2117 (1994).
- [16] T. Zuo, A. D. Bandrauk, and P. B. Corkum, Laser-induced electron diffraction: A new tool for probing ultrafast molecular dynamics, *Chem. Phys. Lett.* **259**, 313 (1996).
- [17] B. Wolter, M. G. Pullen, A. T. Le, M. Baudisch, K. Doblhoff-Dier, A. Senftleben, M. Hemmer, C. D. Schröter, J. Ullrich, T. Pfeifer *et al.*, Ultrafast electron diffraction imaging of bond breaking in di-ionized acetylene, *Science* **354**, 308 (2016).
- [18] S. Ghimire, A. D. DiChiara, E. Sistrunk, P. Agostini, L. F. DiMauro, and D. A. Reis, Observation of high-order harmonic generation in a bulk crystal, *Nat. Phys.* **7**, 138 (2011).
- [19] G. Vampa and T. Brabec, Merge of high harmonic generation from gases and solids and its implications for attosecond science, *J. Phys. B: At. Mol. Opt. Phys.* **50**, 083001 (2017).
- [20] E. N. Osika, A. Chacón, L. Ortmann, N. Suárez, J. A. Pérez-Hernández, B. Szafran, M. F. Ciappina, F. Sols, A. S. Landsman, and M. Lewenstein, Wannier-Bloch Approach to Localization in High-Harmonics Generation in Solids, *Phys. Rev. X* **7**, 021017 (2017).
- [21] G. Vampa, T. J. Hammond, N. Thiré, B. E. Schmidt, F. Légaré, C. R. McDonald, T. Brabec, and P. B. Corkum, Linking high harmonics from gases and solids, *Nature* **522**, 462 (2015).
- [22] G. Vampa, C. R. McDonald, G. Orlando, P. B. Corkum, and T. Brabec, Semiclassical analysis of high harmonic generation in bulk crystals, *Phys. Rev. B* **91**, 064302 (2015).
- [23] G. Vampa, T. J. Hammond, N. Thiré, B. E. Schmidt, F. Légaré, C. R. McDonald, T. Brabec, D. D. Klug, and P. B. Corkum, All-Optical Reconstruction of Crystal Band Structure, *Phys. Rev. Lett.* **115**, 193603 (2015).
- [24] D. Golde, T. Meier, and S. W. Koch, High harmonics generated in semiconductor nanostructures by the coupled dynamics of optical inter- and intraband excitations, *Phys. Rev. B* **77**, 075330 (2008).
- [25] H. Lakhotia, H. Y. Kim, M. Zhan, S. Hu, S. Meng, and E. Goulielmakis, Laser picoscopy of valence electrons in solids, *Nature* **583**, 55 (2020).
- [26] H. Liu, Y. Li, Y. S. You, S. Ghimire, T. F. Heinz, and D. A. Reis, High-harmonic generation from an atomically thin semiconductor, *Nat. Phys.* **13**, 262 (2017).
- [27] T. T. Luu and H. J. Wörner, Measurement of the Berry curvature of solids using high-harmonic spectroscopy, *Nat. Commun.* **9**, 916 (2018).
- [28] H. K. Avetissian and G. F. Mkrtchian, High laser harmonics induced by the Berry curvature in time-reversal invariant materials, [arXiv:2005.09449](https://arxiv.org/abs/2005.09449).
- [29] I. Al-Naib, J. E. Sipe, and M. M. Dignam, High harmonic generation in undoped graphene: Interplay of inter- and intraband dynamics, *Phys. Rev. B* **90**, 245423 (2014).
- [30] D. Dimitrovski, L. B. Madsen, and T. G. Pedersen, High-order harmonic generation from gapped graphene: Perturbative response and transition to nonperturbative regime, *Phys. Rev. B* **95**, 035405 (2017).
- [31] N. Yoshikawa, T. Tamaya, and K. Tanaka, High-harmonic generation in graphene enhanced by elliptically polarized light excitation, *Science* **356**, 736 (2017).
- [32] O. Zurrón, A. Picón, and L. Plaja, Theory of high-order harmonic generation for gapless graphene, *New J. Phys.* **20**, 053033 (2018).
- [33] T. Tamaya, S. Konabe, and S. Kawabata, Orientation dependence of high-harmonic generation in monolayer transition metal dichalcogenides, [arXiv:1706.00548](https://arxiv.org/abs/1706.00548).
- [34] N. Yoshikawa, K. Nagai, K. Uchida, Y. Takaguchi, S. Sasaki, Y. Miyata, and K. Tanaka, Interband resonant high-harmonic generation by valley polarized electron-hole pairs, *Nat. Commun.* **10**, 1 (2019).
- [35] M.-X. Guan, C. Lian, S.-Q. Hu, H. Liu, S.-J. Zhang, J. Zhang, and S. Meng, Cooperative evolution of intraband and interband excitations for high-harmonic generation in strained MoS₂, *Phys. Rev. B* **99**, 184306 (2019).
- [36] L. Jia, Z. Zhang, D. Z. Yang, Y. Liu, M. S. Si, G. P. Zhang, and Y. S. Liu, Optical high-order harmonic generation as a structural characterization tool, *Phys. Rev. B* **101**, 144304 (2020).
- [37] H. K. Avetissian, A. K. Avetissian, A. G. Ghazaryan, G. F. Mkrtchian, and K. V. Sedrakian, High-harmonic generation at particle-hole multiphoton excitation in gapped bilayer graphene, *J. Nanophotonics* **14**, 026004 (2020).
- [38] R. E. F. Silva, I. V. Blinov, A. N. Rubtsov, O. Smirnova, and M. Ivanov, High-harmonic spectroscopy of ultrafast many-body dynamics in strongly correlated systems, *Nat. Photon.* **12**, 266 (2018).
- [39] S. Takayoshi, Y. Murakami, and P. Werner, High-harmonic generation in quantum spin systems, *Phys. Rev. B* **99**, 184303 (2019).
- [40] W. Su, J. Schrieffer, and A. J. Heeger, Solitons in Polyacetylene, *Phys. Rev. Lett.* **42**, 1698 (1979).
- [41] D. Bauer and K. K. Hansen, High-Harmonic Generation in Solids with and without Topological Edge States, *Phys. Rev. Lett.* **120**, 177401 (2018).
- [42] H. Drüeke and D. Bauer, Robustness of topologically sensitive harmonic generation in laser-driven linear chains, *Phys. Rev. A* **99**, 053402 (2019).
- [43] C. Jürß and D. Bauer, High-harmonic generation in Su-Schrieffer-Heeger chains, *Phys. Rev. B* **99**, 195428 (2019).
- [44] R. E. F. Silva, Á. Jiménez-Galán, B. Amorim, O. Smirnova, and M. Ivanov, Topological strong-field physics on sub-laser-cycle timescale, *Nat. Photon.* **13**, 849 (2019).
- [45] S. de Vega, J. D. Cox, F. Sols, and F. J. García de Abajo, Strong-field-driven dynamics and high-harmonic generation in interacting one dimensional systems, *Phys. Rev. Research* **2**, 013313 (2020).
- [46] L. Jia, Z. Zhang, D. Z. Yang, M. S. Si, G. P. Zhang, and Y. S. Liu, High harmonic generation in magnetically-doped topological insulators, *Phys. Rev. B* **100**, 125144 (2019).
- [47] H. K. Avetissian, A. K. Avetissian, B. R. Avchyan, and G. F. Mkrtchian, Multiphoton excitation and high-harmonics generation in topological insulator, *J. Phys.: Condens. Matter* **30**, 185302 (2018).

- [48] C. H. Lee, H. H. Yap, T. Tai, G. Xu, X. Zhang, and J. Gong, Enhanced higher harmonic generation from nodal topology, *Phys. Rev. B* **102**, 035138 (2020).
- [49] L. Asteria, D. T. Tran, T. Ozawa, M. Tarnowski, B. S. Rem, N. Fläschner, K. Sengstock, N. Goldman, and C. Weitenberg, Measuring quantized circular dichroism in ultracold topological matter, *Nat. Phys.* **15**, 449 (2019).
- [50] G. P. Zhang, M. S. Si, M. Murakami, Y. H. Bai, and T. F. George, Generating high-order optical and spin harmonics from ferromagnetic monolayers, *Nat. Commun.* **9**, 1 (2018).
- [51] F. D. M. Haldane, Model for a Quantum Hall Effect Without Landau Levels: Condensed-Matter Realization of the “Parity Anomaly”, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [52] L. Yue and M. B. Gaarde, Structure gauges and laser gauges for the semiconductor Bloch equations in high-order harmonic generation in solids, *Phys. Rev. A* **101**, 053411 (2020).
- [53] M. Kohmoto, Topological invariant and the quantization of the Hall conductance, *Ann. Phys.* **160**, 343 (1985).
- [54] F. Yang and R.-B. Liu, Berry phases of quantum trajectories of optically excited electron-hole pairs in semiconductors under strong terahertz fields, *New J. Phys.* **15**, 115005 (2013).
- [55] N. Saito, P. Xia, F. Lu, T. Kanai, J. Itatani, and N. Ishii, Observation of selection rules for circularly polarized fields in high-harmonic generation from a crystalline solid, *Optica* **4**, 1333 (2017).
- [56] N. R. Cooper, J. Dalibard, and I. B. Spielman, Topological bands for ultracold atoms, *Rev. Mod. Phys.* **91**, 015005 (2019).
- [57] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Quantum spin Hall effect and topological phase transition in HgTe quantum wells, *Science* **314**, 1757 (2006).
- [58] M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, Quantum spin Hall insulator state in HgTe quantum wells, *Science* **318**, 766 (2007).
- [59] C. L. Kane and E. J. Mele, Quantum Spin Hall Effect in Graphene, *Phys. Rev. Lett.* **95**, 226801 (2005).
- [60] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, Experimental realization of the topological Haldane model with ultracold fermions, *Nature* **515**, 237 (2014).
- [61] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, *Rev. Mod. Phys.* **88**, 035005 (2016).
- [62] E. I. Blount, Formalisms of band theory, in *Solid State Physics*, edited by F. Seitz and D. Turnbull (Academic Press, New York, 1962), Vol. 13, pp. 305–373.
- [63] S. Y. Kruchinin, M. Korbman, and V. S. Yakovlev, Theory of strong-field injection and control of photocurrent in dielectrics and wide band gap semiconductors, *Phys. Rev. B* **87**, 115201 (2013).
- [64] M. Spivak, *A Comprehensive Introduction to Differential Geometry* (Publish or Perish, Houston, 1999).
- [65] R. E. F. Silva, F. Martín, and M. Ivanov, High harmonic generation in crystals using maximally localized Wannier functions, *Phys. Rev. B* **100**, 195201 (2019).
- [66] H. Haug and S. W. Koch, *Quantum Theory of the Optical and Electronic Properties of Semiconductors* (World Scientific, Singapore, 2004).
- [67] G. Vampa, C. R. McDonald, G. Orlando, D. D. Klug, P. B. Corkum, and T. Brabec, Theoretical Analysis of High-Harmonic Generation in Solids, *Phys. Rev. Lett.* **113**, 073901 (2014).
- [68] P. Pechukas, T. F. George, K. Morokuma, F. J. McLafferty, and J. R. Laing, Analytic structure of the eigenvalue problem as used in semiclassical theory of electronically inelastic collisions, *J. Chem. Phys.* **64**, 1099 (1976).
- [69] J.-T. Hwang and P. Pechukas, The adiabatic theorem in the complex plane and the semiclassical calculation of nonadiabatic transition amplitudes, *J. Chem. Phys.* **67**, 4640 (1977).
- [70] M. G. Reuter, A unified perspective of complex band structure: Interpretations, formulations, and applications, *J. Phys.: Condens. Matter* **29**, 053001 (2016).
- [71] P. G. Hawkins, M. Y. Ivanov, and V. S. Yakovlev, Effect of multiple conduction bands on high-harmonic emission from dielectrics, *Phys. Rev. A* **91**, 013405 (2015).
- [72] I. Floss, C. Lemell, G. Wachter, V. Smejkal, S. A. Sato, X.-M. Tong, K. Yabana, and J. Burgdörfer, *Ab initio* multiscale simulation of high-order harmonic generation in solids, *Phys. Rev. A* **97**, 011401 (2018).
- [73] A. Fleischer, O. Kfir, T. Diskin, P. Sidorenko, and O. Cohen, Spin angular momentum and tunable polarization in high-harmonic generation, *Nat. Photon.* **8**, 543 (2014).
- [74] O. E. Alon, V. Averbukh, and N. Moiseyev, Selection Rules for the High Harmonic Generation Spectra, *Phys. Rev. Lett.* **80**, 3743 (1998).
- [75] S. A. Sørngård, S. I. Simonsen, and J. P. Hansen, High-order harmonic generation from graphene: Strong attosecond pulses with arbitrary polarization, *Phys. Rev. A* **87**, 053803 (2013).
- [76] Z.-Y. Chen and R. Qin, Circularly polarized extreme ultraviolet high harmonic generation in graphene, *Opt. Express* **27**, 3761 (2019).
- [77] Ó. Zurrón-Cifuentes, R. Boyero-García, C. Hernández-García, A. Picón, and L. Plaja, Optical anisotropy of non-perturbative high-order harmonic generation in gapless graphene, *Opt. Express* **27**, 7776 (2019).
- [78] Á. Jiménez-Galán, R. E. F. Silva, O. Smirnova, and M. Ivanov, Ultrafast topology for strong-field valleytronics, [arXiv:1910.07398](https://arxiv.org/abs/1910.07398).
- [79] F. Mauger, A. D. Bandrauk, and T. Uzer, Circularly polarized molecular high harmonic generation using a bicircular laser, *J. Phys. B: At. Mol. Opt. Phys.* **49**, 10LT01 (2016).
- [80] D. Baykusheva, Md. S. Ahsan, N. Lin, and H. J. Wörner, Bicircular High-Harmonic Spectroscopy Reveals Dynamical Symmetries of Atoms and Molecules, *Phys. Rev. Lett.* **116**, 123001 (2016).
- [81] K.-J. Yuan and A. D. Bandrauk, Symmetry in circularly polarized molecular high-order harmonic generation with intense bicircular laser pulses, *Phys. Rev. A* **97**, 023408 (2018).
- [82] K. Konishi, T. Higuchi, J. Li, J. Larsson, S. Ishii, and M. Kuwata-Gonokami, Polarization-Controlled Circular Second-Harmonic Generation from Metal Hole Arrays with Threefold Rotational Symmetry, *Phys. Rev. Lett.* **112**, 135502 (2014).
- [83] L. Medišauskas, J. Wragg, H. van der Hart, and M. Yu. Ivanov, Generating Isolated Elliptically Polarized Attosecond Pulses using Bichromatic Counterrotating Circularly Polarized Laser Fields, *Phys. Rev. Lett.* **115**, 153001 (2015).
- [84] E. Pisanty and A. Jiménez-Galán, Strong-field approximation in a rotating frame: High-order harmonic emission from p states in bicircular fields, *Phys. Rev. A* **96**, 063401 (2017).

- [85] Y. Harada, E. Haraguchi, K. Kaneshima, and T. Sekikawa, Circular dichroism in high-order harmonic generation from chiral molecules, *Phys. Rev. A* **98**, 021401 (2018).
- [86] D. Ayuso, P. Decleva, S. Patchkovskii, and O. Smirnova, Chiral dichroism in bi-elliptical high-order harmonic generation, *J. Phys. B: At. Mol. Opt. Phys.* **51**, 06LT01 (2018).
- [87] K. G. Wilson, Quarks and strings on a lattice, in *New Phenomena in Subnuclear Physics*, edited by A. Zichichi (Springer US, Boston, Mass, 1977), pp. 69–142.
- [88] J. B. Kogut, The lattice gauge theory approach to quantum chromodynamics, *Rev. Mod. Phys.* **55**, 775 (1983).
- [89] L. H. Karsten and J. Smith, Lattice fermions: Species doubling, chiral invariance and the triangle anomaly, *Nucl. Phys. B* **183**, 103 (1981).
- [90] H. B. Nielsen and M. Ninomiya, Absence of neutrinos on a lattice: (I). Proof by homotopy theory, *Nucl. Phys. B* **185**, 20 (1981).
- [91] F. Wilczek, Problem of Strong P and T Invariance in the Presence of Instantons, *Phys. Rev. Lett.* **40**, 279 (1978).
- [92] S. Weinberg, A New Light Boson? *Phys. Rev. Lett.* **40**, 223 (1978).
- [93] J. Preskill, M. B. Wise, and F. Wilczek, Cosmology of the invisible axion, *Phys. Lett. B* **120**, 127 (1983).
- [94] X.-L. Qi, T. L. Hughes, and S.-C. Zhang, Topological field theory of time-reversal invariant insulators, *Phys. Rev. B* **78**, 195424 (2008).
- [95] A. M. Essin, J. E. Moore, and D. Vanderbilt, Magnetoelectric Polarizability and Axion Electrodynamics in Crystalline Insulators, *Phys. Rev. Lett.* **102**, 146805 (2009).
- [96] X.-L. Qi, R. Li, J. Zang, and S.-C. Zhang, Inducing a magnetic monopole with topological surface states, *Science* **323**, 1184 (2009).
- [97] G. Rosenberg and M. Franz, Witten effect in a crystalline topological insulator, *Phys. Rev. B* **82**, 035105 (2010).
- [98] A. Bermudez, L. Mazza, M. Rizzi, N. Goldman, M. Lewenstein, and M. A. Martin-Delgado, Wilson Fermions and Axion Electrodynamics in Optical Lattices, *Phys. Rev. Lett.* **105**, 190404 (2010).
- [99] P. Rodriguez-Lopez, W. J. M. Kort-Kamp, D. A. R. Dalvit, and L. M. Woods, Nonlocal optical response in topological phase transitions in the graphene family, *Phys. Rev. Mater.* **2**, 014003 (2018).
- [100] M. Neupane, A. Richardella, J. Sánchez-Barriga, S. Y. Xu, N. Alidoust, I. Belopolski, C. Liu, G. Bian, D. Zhang, D. Marchenko *et al.*, Observation of quantum-tunnelling-modulated spin texture in ultrathin topological insulator Bi₂Se₃ films, *Nat. Commun.* **5**, 3841 (2014).
- [101] Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, Unconventional superconductivity in magic-angle graphene superlattices, *Nature* **556**, 43 (2018).
- [102] Y. Cao, V. Fatemi, A. Demir, S. Fang, S. L. Tomarken, J. Y. Luo, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras *et al.*, Correlated insulator behaviour at half-filling in magic-angle graphene superlattices, *Nature* **556**, 80 (2018).
- [103] X. Lu, P. Stepanov, W. Yang, M. Xie, M. A. Aamir, I. Das, C. Urgell, K. Watanabe, T. Taniguchi, G. Zhang *et al.*, Superconductors, orbital magnets and correlated states in magic-angle bilayer graphene, *Nature* **574**, 653 (2019).
- [104] L. Balents, C. R. Dean, D. K. Efetov, and A. F. Young, Superconductivity and strong correlations in moiré flat bands, *Nat. Phys.* **16**, 725 (2020).
- [105] A. Acín, I. Bloch, H. Buhrman, T. Calarco, C. Eichler, J. Eisert, D. Esteve, N. Gisin, S. J. Glaser, F. Jelezko *et al.*, The quantum technologies roadmap: a European community view, *New J. Phys.* **20**, 080201 (2018).
- [106] M. Lewenstein, A. Sanpera, and V. Ahufinger, *Ultracold Atoms in Optical Lattices: Simulating Quantum Many-Body Systems* (Oxford University Press, 2012).
- [107] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with ultracold gases, *Rev. Mod. Phys.* **80**, 885 (2008).
- [108] J. I. Cirac and P. Zoller, Goals and opportunities in quantum simulation, *Nat. Phys.* **8**, 264 (2012).
- [109] I. Bloch, J. Dalibard, and S. Nascimbène, Quantum simulations with ultracold quantum gases, *Nat. Phys.* **8**, 267 (2012).
- [110] R. Blatt and C. F. Roos, Quantum simulations with trapped ions, *Nat. Phys.* **8**, 277 (2012).
- [111] A. Aspuru-Guzik and P. Walther, Photonic quantum simulators, *Nat. Phys.* **8**, 285 (2012).
- [112] A. A. Houck, H. E. Türeci, and J. Koch, On-chip quantum simulation with superconducting circuits, *Nat. Phys.* **8**, 292 (2012).
- [113] S. Sala, J. Förster, and A. Saenz, Ultracold-atom quantum simulator for attosecond science, *Phys. Rev. A* **95**, 011403 (2017).
- [114] R. Senaratne, S. V. Rajagopal, T. Shimasaki, P. E. Dotti, K. M. Fujiwara, K. Singh, Z. A. Geiger, and D. M. Weld, Quantum simulation of ultrafast dynamics using trapped ultracold atoms, *Nat. Commun.* **9**, 2065 (2018).
- [115] R. Ramos, D. Spierings, I. Racicot, and A. M. Steinberg, Measurement of the time spent by a tunnelling atom within the barrier region, *Nature* **583**, 529 (2020).
- [116] <https://github.com/ftachacon/NonLinearCircularDichroism>.